

Poster Presentation Number 17, Session 1, Thursday 17:45-19:15

First insights into trabecular bone distribution in the first metacarpal of *Homo naledi*

Christopher J. Dunmore¹, Sebastian Bachmann², Alexander Synek², Tracy L. Kivell^{3,4}, Matthew M. Skinner^{1,3,4}

1 - Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, UK · 2 - Institute of Lightweight Design and Structural Biomechanics, TU Wien, Vienna, Austria · 3 - Department of Human Origins, Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany · 4 - Centre for the Exploration of the Deep Human Journey, University of the Witwatersrand, Johannesburg, South Africa

Modern human dexterity is thought to be facilitated by our distinctive pollical (thumb) morphology. The shape of the first metacarpal (Mc1), including the robusticity of the shaft and disto-radial flange as well as the articular surfaces of the first metacarpophalangeal and trapeziometacarpal joints, have been used to infer the degree of dexterity in several hominin species [1-4]. The *Homo naledi* Mc1 has a unique external morphology among known fossil hominins with no direct analogue in extant primates, possessing a small proximal articular surface, a flat radio-palmarly extended distal articular surface, and both radial as well as ulnar diaphyseal flanges [3-4]. As trabecular bone can (re)model in response to mechanical loading over an individual's lifetime and is located beneath both epiphyseal and diaphyseal Mc1 external morphology, it may provide additional functional information about *H. naledi* thumb use. Compared to other living great apes, humans have significantly more trabecular bone directly beneath the radial aspect of both the distal and proximal articulations of the first metacarpal, as well as beneath the disto-radial flange. These differences are consistent with differences in both external Mc1 morphology and observed manipulative capabilities [5].

Here, we use canonical holistic morphometric analysis (cHMA) to statistically compare the distribution of relative trabecular bone density (rBV/TV) in largely complete *H. naledi* Mc1s (U.W. 101-1321 and U.W. 101-270) to a sample of living great apes including: modern humans (*Homo sapiens*, n=10), bonobos (*Pan paniscus*, n=10), chimpanzees (*Pan troglodytes*, n=11), gorillas (*Gorilla gorilla*, n=10) and orangutans (*Pongo* sp., n=7). Run in Medtool 4.5 (www.medtool.at), this method uses a statistical deformation model to create a canonical Mc1, to which all Mc1s are subsequently deformed. This controls for shape differences before statistical volumetric comparison of trabecular distribution. The trabecular distribution of the whole bone, the proximal epiphysis, the distal epiphysis, and the diaphyseal medullary cavity of the fossils, were each separately compared to the homologous region of the comparative sample.

H. naledi Mc1s demonstrate some intraspecific variation in trabecular distribution. We find evidence for radial concentration of trabecular bone beneath the proximal and distal articular surfaces of U.W. 101-1321 consistent with the modern human pattern, thought to reflect thumb loading during forceful opposition. U.W. 101-270 does not show this signal as clearly. However, both fossils show trabecular concentrations beneath both the ulnar and radial diaphyseal flanges, suggesting a plastic response to strong first dorsal interosseous and opponens pollicis muscle action. We hypothesize that the antagonistic actions of these muscles assisted in the maintenance of stability at the small proximal articulation of the Mc1 in *H. naledi*, but ongoing analysis of the large hypodigm available for this species is needed to properly support this supposition.

We thank the following researchers or curators for access to specimens in their care: I. Livne and R. Jennings (Powell-Cotton Museum), A. vanHeteren and M. Hiermeier (Zoologische Staatssammlung München), C. Boesch and U. Schwarz (Max Planck Institute for Evolutionary Anthropology), A. Ragni (Smithsonian Institution, National Museum of Natural History); M. Teschler-Nicola and R. Muehl (Natural History Museum, Vienna), J. Moggi-Cecchi and S. Bortoluzzi (University of Florence), F. Mayer (Museum für Naturkunde—Leibniz Institute for Evolution and Biodiversity Science, Berlin), B. Großkopf (Johann-Friedrich-Blumenbach-Institut für Zoologie und Anthropologie der Georg-August-Universität Göttingen), E. Gilissen and W. Wendelen (Musée Royal de l'Afrique Centrale), V. Volpato (Senckenberg Museum of Frankfurt), B. Zipfel and L. Berger (University of the Witwatersrand), E. Gilissen and W. Wendelen (Musée Royal de l'Afrique Centrale), V. Volpato (Senckenberg Museum of Frankfurt). This research was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant no. 819960).

References-[1] Galletta L., Stephens, N.B., Bardo, A., Kivell, T.L., Marchi, D., 2019. Three-dimensional geometric morphometric analysis of the first metacarpal distal articular surface in humans, great apes and fossil hominins. *Journal of Human Evolution* 132, 119-136. [2] Marchi, D., Proctor, D.J., Huston, E., Nicholas, C.L., Fischer, F., 2017. Morphological correlates of the first metacarpal proximal articular surface with manipulative capabilities in apes, humans and South African early hominins. *Comptes Rendus Palevol* 16(5-6), 645-654. [3] Bowland, L.A., Scott, J.E., Kivell, T.L., Patel, B.A., Tocheri, M.W., Orr, C.M., 2021. *Homo naledi* pollical metacarpal shaft morphology is distinctive and intermediate between that of australopiths and other members of the genus *Homo*. *Journal of Human Evolution* 158, 103048. [4] Kivell, T., Deane, A., Tocheri, M. W., Orr, C.M., Schmid, P., Hawks, J., Berger, L.R., Churchill, S.E., 2015. The hand of *Homo naledi*. *Nature Communications* 6, 8431. [5] Dunmore, C.J., Bachmann, S., Synek, A., Pahr, D.H., Skinner, M.M., Kivell, T.L., in press. The deep trabecular structure of first metacarpals in extant hominids. *American Journal of Biological Anthropology*.