



# Global Existence and Weak-Strong Uniqueness for Chemotaxis Compressible Navier–Stokes Equations Modeling Vascular Network Formation

Xiaokai Huo  and Ansgar Jüngel

Communicated by E. Feireisl

**Abstract.** A model of vascular network formation is analyzed in a bounded domain, consisting of the compressible Navier–Stokes equations for the density of the endothelial cells and their velocity, coupled to a reaction-diffusion equation for the concentration of the chemoattractant, which triggers the migration of the endothelial cells and the blood vessel formation. The coupling of the equations is realized by the chemotaxis force in the momentum balance equation. The global existence of finite energy weak solutions is shown for adiabatic pressure coefficients  $\gamma > 8/5$ . The solutions satisfy a relative energy inequality, which allows for the proof of the weak–strong uniqueness property.

**Mathematics Subject Classification.** 35Q30, 35K57, 35K65, 76N05.

**Keywords.** Compressible Navier–Stokes equations, Chemotaxis force, Global existence of solutions, Weak–strong uniqueness, Relative energy.

## 1. Introduction

The formation of blood vessels is regulated by chemical signals triggering the movement of endothelial cells. The cells may self-assemble into a vascular network, which is known as vasculogenesis. In this paper, we analyze a mathematical model for the formation of vascular networks, based on mass and momentum balance equations including a chemotaxis force and coupled with a reaction-diffusion equation for the signal concentration. The existence of global weak solutions to the resulting chemotaxis compressible Navier–Stokes equations was proved in [1] for pressures with adiabatic exponent  $\gamma > 3$ . We extend the existence result to the range  $\gamma > 8/5$  and prove a weak–strong uniqueness property. The proofs are based on a new relative energy inequality.

The dynamics of the density  $\rho(x, t)$  of the endothelial cells, their velocity  $v(x, t)$ , and the concentration  $c(x, t)$  of the chemoattractant (e.g. the vascular endothelial growth factor VEGF-A [17]) is given by the equations

$$\partial_t \rho + \operatorname{div}(\rho v) = 0, \tag{1}$$

$$\partial_t(\rho v) + \operatorname{div}(\rho v \otimes v) + \nabla p(\rho) = \mu \Delta v + (\lambda + \mu) \nabla \operatorname{div} v + \rho \nabla c - \frac{\rho v}{\zeta}, \tag{2}$$

$$\partial_t c = \Delta c - c + \rho \quad \text{in } \Omega, t > 0, \tag{3}$$

where  $\Omega \subset \mathbb{R}^3$  is a bounded domain,  $p(\rho) = \rho^\gamma$  with the adiabatic exponent  $\gamma > 1$  is the pressure, the Lamé viscosity constants  $\mu, \lambda$  satisfy  $\mu > 0$  and  $3\lambda + 2\mu > 0$ , and  $\zeta > 0$  is a relaxation constant. We

---

The last author acknowledges partial support from the Austrian Science Fund (FWF), Grants P33010 and F65. This work has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme, ERC Advanced Grant no. 101018153.

impose the initial and boundary conditions

$$\rho(\cdot, 0) = \rho^0, \quad v(\cdot, 0) = v^0, \quad c(\cdot, 0) = c^0 \quad \text{in } \Omega, \tag{4}$$

$$v = 0, \quad \nabla c \cdot \nu = 0 \quad \text{on } \partial\Omega, \quad t > 0. \tag{5}$$

The boundary condition for the velocity  $v$  is the no-slip condition, and the no-flux boundary condition for  $c$  means that there is no inflow or outflow of the concentration. The momentum balance equation (2) includes viscous terms as in [1] (suggested in [2, p. 1862]) as well as the chemotaxis force  $\rho f_{\text{chem}} = -\rho \nabla c$  and the drag force  $\rho f_{\text{drag}} = -\rho v / \zeta$ . The reaction-diffusion equation (3) for the signal concentration models diffusion in the surrounding medium, degradation of the signal in finite time, and the release of the signal produced by the cells. We have set the physical constants in (1)–(3) equal to one, except  $\zeta$  to distinguish terms originating from the drag force.

The existence of global finite energy weak solutions to (1)–(5) has been proved in [1] for  $\gamma > 3$ . This restriction comes from the estimation of the chemotaxis force; see Remark 4 on page 10. We extend the existence result to  $\gamma > 8/5$  by rewriting the force term  $\rho \nabla c$  via (3) as  $(\partial_t c - \Delta c + c) \nabla c$  and exploiting the properties of the Bogovskii operator. Replacing the parabolic equation (3) for  $c$  by the elliptic one, we can even allow for  $\gamma > 3/2$ , which is the condition needed in the existence theory of the compressible Navier–Stokes equations [8]; see Remark 6. This may indicate that our condition  $\gamma > 8/5$  for system (1)–(3) is not optimal. We discuss this issue further in Remark 5.

The idea of the existence proof in [1] is to derive a priori estimate via the energy-type functional

$$\tilde{H}(\rho, v, c) = \int_{\Omega} \left( \psi(\rho) + \frac{1}{2} \rho |v|^2 + \frac{1}{2} c^2 \right) dx,$$

where  $\psi(\rho) = \rho \int_0^\rho s^{-2} p(s) ds = \rho^\gamma / (\gamma - 1)$  can be interpreted as the internal energy. Unfortunately, this functional is not bounded as  $t \rightarrow \infty$ . Our idea is to use the physical (free) energy,

$$E(\rho, v, c) = \int_{\Omega} \left( \psi(\rho) + \frac{1}{2} \rho |v|^2 + \frac{1}{2} (|\nabla c|^2 + c^2) - \rho c \right) dx, \tag{6}$$

which is the sum of the kinetic energy  $\frac{1}{2} \int_{\Omega} \rho |v|^2 dx$  and the energy  $E(\rho, 0, c)$  of the parabolic–parabolic Keller–Segel model. We show in Sect. 2 (see Lemma 3 on page 6) that

$$\frac{dE}{dt}(\rho, v, c) + \int_{\Omega} (\mu |\nabla v|^2 + (\lambda + \mu) |\operatorname{div} v|^2) dx + \int_{\Omega} |\partial_t c|^2 dx \leq 0,$$

providing a bound for  $E((\rho, v, c)(t))$  uniformly in time. Clearly, to infer a priori estimates, we need an upper bound for  $\rho c$ . This is done by using the inequality

$$\int_{\Omega} \rho c dx \leq \frac{1}{2} \|\psi(\rho)\|_{L^1(\Omega)} + \frac{1}{4} \|\nabla c\|_{L^2(\Omega)}^2 + C_1(\gamma) \|c\|_{L^1(\Omega)}^{C_2(\gamma)}, \tag{7}$$

which is due to Sugiyama [18] (see Lemma 9 on page 21), where  $C_1(\gamma) > 0$  and  $C_2(\gamma) > 0$  only depend on  $\gamma$ , and which (7) requires the condition  $\gamma > 8/5$ . The first two terms on the right-hand side of (7) can be absorbed by the energy, while the  $L^1(\Omega)$  norm of  $c$  can be bounded in terms of the initial data  $(\rho^0, c^0)$ . This provides a bound for the modified energy-type functional

$$H(\rho, v, c) = \frac{1}{2} \int_{\Omega} \left( \psi(\rho) + \rho |v|^2 + \frac{1}{2} |\nabla c|^2 + c^2 \right) dx, \tag{8}$$

namely

$$\begin{aligned} H((\rho, v, c)(t)) &+ \int_0^t \int_{\Omega} (\mu |\nabla v|^2 + (\lambda + \mu) |\operatorname{div} v|^2) dx ds \\ &+ \int_0^t \int_{\Omega} |\partial_s c|^2 dx ds \leq C(\rho^0, v^0, c^0), \end{aligned}$$

which allows us to prove the global existence of finite energy weak solutions such that  $H(\rho, v, c)$  is finite for all  $t > 0$ . This type of solutions is defined as follows.

**Definition 1** (*Finite energy weak solution*). The triple  $(\rho, v, c)$  is a *finite energy weak solution* to (1)–(5) if

- they satisfy the regularity

$$\begin{aligned} \rho &\in L^\infty(0, T; L^\gamma(\Omega)), \quad \rho \geq 0 \text{ in } \Omega, \quad t > 0, \\ v &\in L^2(0, T; H_0^1(\Omega; \mathbb{R}^3)), \quad c \in L^\infty(0, T; H^1(\Omega)) \cap H^1(0, T; L^2(\Omega)); \end{aligned}$$

- equation (1) is satisfied in the sense of renormalized solutions [6, Section 10.18];
- equations (2)–(3) are satisfied in the sense of distributions;
- the energy inequality

$$\begin{aligned} E((\rho, v, c)(t)) &+ \int_0^t \int_\Omega (\mu |\nabla v|^2 + (\lambda + \mu) |\operatorname{div} v|^2) dx ds \\ &+ \int_0^t \int_\Omega |\partial_s c_n|^2 dx ds + \frac{1}{\zeta} \int_0^t \int_\Omega \rho_n |v_n|^2 dx ds \leq E(\rho^0, v^0, c^0) \end{aligned}$$

holds for a.e.  $t \in (0, T)$ .

We introduce for  $1 < p, q < \infty$  the space  $W_\nu^{2-2/p, q}(\Omega)$  as the completion of the space of functions  $w \in C^\infty(\bar{\Omega})$  satisfying  $\nabla w \cdot \nu = 0$  on  $\partial\Omega$  in the norm of  $W^{2-2/p, q}(\Omega)$ . We can now state our first main result.

**Theorem 1** (*Global existence*). Let  $\partial\Omega \in C^2$ ,  $p(\rho) = \rho^\gamma$  for  $\rho \geq 0$  with  $\gamma > 8/5$ . Assume that the initial datum satisfies  $\rho^0 \in L^\gamma(\Omega)$  with  $\rho^0 \geq 0$ ,  $\rho^0 \not\equiv 0$  in  $\Omega$ ,  $\rho^0 |v^0|^2 \in L^1(\Omega)$ , and  $c^0 \in W_\nu^{2-2/\gamma, \gamma}(\Omega)$ ,  $c^0 \geq 0$  in  $\Omega$ . Then there exists a finite energy weak solution  $(\rho, v, c)$  to (1)–(5) in the sense of Definition 1.

The condition on the initial datum  $c^0 \in W_\nu^{2-2/\gamma, \gamma}(\Omega)$  can be rephrased in terms of interpolation or Besov spaces. Indeed, the condition is needed to apply the maximal regularity result of Theorem 10, and the regularity on the initial datum can be formulated in such spaces; see [6, Theorem 10.22]. The definition of the pressure can be relaxed to  $p \in C^0([0, \infty)) \cap C^2(0, \infty)$ ,  $p(0) = 0$ ,  $p'(\rho) > 0$  for  $\rho > 0$ , and  $\rho^{1-\gamma} p'(\rho) \rightarrow a > 0$  as  $\rho \rightarrow \infty$ ; see [7, (2.1)]. The proof of the theorem is based on the existence theory for the compressible Navier–Stokes equations [8]. More precisely, we add some artificial diffusion and an artificial pressure term, construct Faedo–Galerkin solutions to the approximate problem, prove an approximate energy inequality for these solutions, and pass to the de-regularizing limit. Improved uniform bounds for the cell density in  $L^{\gamma+\theta}(\Omega)$  for some  $\theta > 0$  are derived by testing the mass balance equation with a test function involving the Bogovskii operator. The novel part is the estimate of the chemotaxis force term.

Next, we formulate the weak–strong uniqueness property of the system, meaning that a weak and a strong solution emanating from the same initial data coincide as long as the latter exists.

**Theorem 2** (*Weak–strong uniqueness*). Let  $(\rho, v, c)$  and  $(\bar{\rho}, \bar{v}, \bar{c})$  be two finite energy weak solutions to (1)–(5) constructed in Theorem 1 with the same initial data. Assume that  $(\bar{\rho}, \bar{v}, \bar{c})$  satisfies the additional regularity

$$0 < c_p \leq \bar{\rho} \leq C_p, \quad |\bar{v}| \leq C_v \text{ a.e. in } \Omega \times (0, T), \quad |\nabla \bar{\rho}|, |\nabla^2 \bar{v}| \in L^2(0, T; L^q(\Omega)) \tag{9}$$

for  $q > 3$  and some constants  $c_p, C_p, C_v > 0$ . Then  $\rho = \bar{\rho}$ ,  $v = \bar{v}$ , and  $c = \bar{c}$  in  $\Omega \times (0, T)$ .

The no-vacuum assumption  $\bar{\rho} \geq c_p > 0$  was also needed in [9] and in related contexts, e.g. for the weak–strong uniqueness property of Maxwell–Stefan systems [11]. The proof of Theorem 2 is based on the relative energy method. The relative energy, associated to the energy functional (6), is given by

$$E(\rho, v, c | r, u, z) = \int_\Omega \left( \psi(\rho | r) + \frac{1}{2} \rho |v - u|^2 + \frac{1}{2} (|\nabla(c - z)|^2 + (c - z)^2) - (\rho - r)(c - z) \right) dx,$$

where  $\psi(\rho|r) = \psi(\rho) - \psi(r) - \psi'(r)(\rho - r)$  is the Bregman distance associated to  $\psi$ . We show in Lemma 7 on page 11 that

$$\begin{aligned}
 E((\rho, v, c)(t)|(r, u, z)(t)) &+ \int_0^t \int_{\Omega} (\mu|\nabla(v - u)|^2 + (\lambda + \mu)|\operatorname{div}(v - u)|^2) dx ds \\
 &+ \int_0^t \int_{\Omega} |\partial_s(c - z)|^2 dx ds \leq E(\rho^0, v^0, c^0|r^0, u^0, z^0) + \int_0^t R(\rho, v, c|r, u, z) ds,
 \end{aligned} \tag{10}$$

where  $(\rho, v, c)$  is a finite energy weak solution to (1)–(5),  $(r, u, z)$  are smooth functions, and the remainder  $R(\rho, v, c|r, u, z)$  is defined in Lemma 7 below. Finite energy weak solutions to the compressible Navier–Stokes equations satisfying (10) have been called *suitable weak solutions* in [9]. It was shown in [7] that finite energy weak solutions in fact always satisfy the relative energy inequality (10) for smooth functions  $(r, u, z)$ .

Defining the modified relative energy

$$H(\rho, v, c|r, u, z) = \frac{1}{2} \int_{\Omega} \left( \psi(\rho|r) + \rho|v - u|^2 + \frac{1}{2}|\nabla(c - z)|^2 + (c - z)^2 \right) dx$$

and giving another weak solution  $(r, u, z) = (\bar{\rho}, \bar{v}, \bar{c})$  satisfying the regularity (9), the idea of the proof is to show that

$$\begin{aligned}
 R(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) &\leq CH(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) \quad \text{and} \\
 \int_{\Omega} (\rho - \bar{\rho})(c - \bar{c}) dx &\leq \frac{1}{2}H(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) + C(\rho^0, c^0),
 \end{aligned}$$

which leads to

$$\frac{1}{2}H((\rho, v, c)(t)|(\bar{\rho}, \bar{v}, \bar{c})(t)) \leq E(\rho^0, v^0, c^0|r^0, u^0, z^0) + C(\rho^0, c^0) + C \int_0^t H(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) ds$$

and which implies, by Gronwall’s lemma, that  $H((\rho, v, c)(t)|(\bar{\rho}, \bar{v}, \bar{c})(t)) = 0$ . Consequently  $\rho(t) = \bar{\rho}(t)$ ,  $v(t) = \bar{v}(t)$ , and  $c(t) = \bar{c}(t)$  for  $t > 0$ .

We finish the introduction by discussing the state of the art. The global existence of finite energy weak solutions to the compressible Navier–Stokes equations with adiabatic exponents  $\gamma > 3/2$  was shown in [8]. The range of  $\gamma$  can be extended to  $\gamma > 1$  for axisymmetric initial data [12] or for a class of density-dependent viscosity coefficients [14], for instance. Germain [10] proved a relative energy inequality and established the weak–strong uniqueness property for solutions to the compressible Navier–Stokes equations with an integrable spatial density gradient. Feireisl et al. [9] proved the existence of so-called suitable weak solutions satisfying a general relative energy inequality with respect to any sufficiently regular pair of functions and concluded the weak–strong uniqueness property.

Compressible Euler equations with chemotaxis force have been introduced in [17] to describe early stages of vasculogenesis. As remarked in [2, Section 3], the fluid equations may also include viscous terms. This leads to chemotaxis compressible Navier–Stokes equations, which have been analyzed in [1] with the pressure function  $p(\rho) = \max\{0, \rho - \rho_c\}^\gamma$ , where  $\gamma > 3$  and  $\rho_c > 0$  is the so-called close-packing density. A viscoelastic mechanical interaction of the cells with the substratum was added to the compressible Euler equations in [20]. Related models are the incompressible Navier–Stokes equations coupled to the chemotaxis Keller–Segel system via the fluid velocity, proposed in [21] and analyzed in, e.g., [22].

The paper is organized as follows. Section 2 is devoted to the proof of Theorem 1. The technical relative energy inequality (10) is proved in Sect. 3. Based on this inequality, Theorem 2 is then shown in Sect. 4. Finally, some auxiliary results are presented in Appendix 4.

## 2. Global Existence of Solutions

In this section, we prove Theorem 1. For this, we proceed as in [8] by constructing an approximate scheme based on a regularized system, deriving uniform energy estimates, and assing to the de-regularization

limit. For later use, we note the relations between the pressure  $p(\rho)$  and the associated internal energy  $\psi(\rho) = \rho \int_0^\rho s^{-2} p(s) ds$ :

$$p(\rho) = \rho\psi'(\rho) - \psi(\rho), \quad \nabla p(\rho) = \rho\nabla\psi'(\rho) \quad \text{for smooth } \rho. \tag{11}$$

### 2.1. Regularized System

We solve first the following regularized system for  $\delta > 0$ ,  $\varepsilon > 0$ , and  $\beta > 4$ :

$$\partial_t \rho + \operatorname{div}(\rho v) = \varepsilon \Delta \rho, \quad \partial_t c = \Delta c - c + \rho, \tag{12}$$

$$\begin{aligned} \partial_t(\rho v) + \operatorname{div}(\rho v \otimes v) + \nabla p(\rho) + \varepsilon \nabla \rho \cdot \nabla v + \delta \nabla \rho^\beta \\ = \mu \Delta v + (\lambda + \mu) \nabla \operatorname{div} v + \rho \nabla c - \frac{\rho v}{\zeta} \quad \text{in } \Omega, \quad t > 0, \end{aligned} \tag{13}$$

subject to the initial and boundary conditions

$$\rho(\cdot, 0) = \rho_\delta^0, \quad v(\cdot, 0) = v^0, \quad c(\cdot, 0) = c^0 \quad \text{in } \Omega, \tag{14}$$

$$\nabla \rho \cdot \nu = 0, \quad v = 0, \quad \nabla c \cdot \nu = 0 \quad \text{on } \partial\Omega, \quad t > 0, \tag{15}$$

where  $\rho_\delta^0$  is a smooth strictly positive function such that  $\rho_\delta^0 \rightarrow \rho^0$  strongly in  $L^\gamma(\Omega)$ . The artificial viscosity term  $\varepsilon \Delta \rho$  is balanced by the term  $\varepsilon \nabla \rho \cdot \nabla v$  in the momentum equation to control the energy. The artificial pressure term  $\delta \nabla \rho^\beta$  is needed to derive an  $L^{\gamma+\theta}(\Omega)$  estimate for the density with  $\theta > 0$ .

The existence of strong solutions to (12)–(15) was shown in [8, Section 2] without the chemotaxis term  $\rho \nabla c$ . Here, we sketch the proof for the problem including the chemotaxis coupling. As in [8], we use the Faedo–Galerkin method. Let  $(\psi_n)$  be a sequence of eigenfunctions of the Laplacian with homogeneous Dirichlet boundary conditions and let  $X_n = \operatorname{span}\{\psi_1, \dots, \psi_n\}$ . Then, following the proof of [16, Section 7.7] or [5, Chapter 7], we obtain the existence of a unique local strong solution  $(\rho_n, v_n, c_n)$  on  $(0, T_n)$  such that  $v_n \in C^1([0, T_n]; X_n)$  and

$$\begin{aligned} \rho_n, \partial_t \rho_n, \nabla \rho_n, \nabla^2 \rho_n, c_n, \partial_t c_n, \nabla c_n, \nabla^2 c_n \quad \text{are Hölder continuous on } \overline{\Omega} \times [0, T_n], \\ \rho_n(x, t) > 0, \quad c_n(x, t) \geq 0 \quad \text{for any } (x, t) \in \overline{\Omega} \times [0, T_n]. \end{aligned}$$

To obtain global solutions, i.e.  $T = T_n$ , we derive an energy inequality for the approximate system.

### 2.2. Energy Inequality for the Approximate System

An energy-type inequality has been derived in [1, Section 2.2]. Here, we use a different energy functional by including the  $H^1(\Omega)$  norm of  $c$ . We show an inequality for the energies  $E(\rho, v, c)$  and  $H(\rho, v, c)$ , defined in (6) and (8), respectively.

**Lemma 3.** *Let  $(\rho_n, v_n, c_n)$  be a strong solution to (12)–(15) constructed in the previous subsection. Then there exists  $C > 0$  independent of  $(n, \delta, \varepsilon)$  such that for any  $0 < t < T_n$ ,*

$$\begin{aligned} E((\rho_n, v_n, c_n)(t)) + \int_0^t \int_\Omega (\mu |\nabla v_n|^2 + (\lambda + \mu) |\operatorname{div} v_n|^2) dx ds + \frac{4\delta\varepsilon}{\beta} \int_0^t \int_\Omega |\nabla \rho_n^{\beta/2}|^2 dx ds \\ + \frac{4\varepsilon}{\gamma} \int_0^t \int_\Omega |\nabla \rho_n^{\gamma/2}|^2 dx ds + \left(1 - \frac{\varepsilon}{4}\right) \int_0^t \int_\Omega |\partial_s c_n|^2 dx ds + \frac{1}{\zeta} \int_0^t \int_\Omega \rho_n |v_n|^2 dx ds \\ \leq E(\rho^0, v^0, v^0) + 2\varepsilon \int_\Omega \rho^\gamma dx + C\varepsilon, \\ H((\rho_n, v_n, c_n)(t)) + \int_0^t \int_\Omega (\mu |\nabla v_n|^2 + (\lambda + \mu) |\operatorname{div} v_n|^2) dx ds \\ + \frac{4\delta\varepsilon}{\beta} \int_0^t \int_\Omega |\nabla \rho_n^{\beta/2}|^2 dx ds + \left(1 - \frac{\varepsilon}{4}\right) \int_\Omega \int_\Omega |\partial_s c_n|^2 dx ds + \frac{1}{\zeta} \int_0^t \int_\Omega \rho_n |v_n|^2 dx ds \end{aligned}$$

$$\leq (E(\rho_\delta^0, v^0, c^0) + C(\rho^0, v^0) + C\varepsilon t)e^{C\varepsilon t}.$$

*Proof. Step 1: Energy inequality for E.* We choose the test function  $\psi'(\rho_n) - \frac{1}{2}|v_n|^2 + \delta\beta\rho_n^{\beta-1}/(\beta - 1)$  in the weak formulation of the first equation in (12) and the test function  $v_n$  in the weak formulation of (13). Adding both equations and taking into account (11), some terms cancel, and we arrive after a standard computation at

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \left( \psi(\rho_n) + \frac{1}{2}\rho_n|v_n|^2 + \frac{\delta}{\beta-1}\rho_n^\beta \right) dx + \int_{\Omega} (\mu|\nabla v_n|^2 + (\lambda + \mu)|\operatorname{div} v_n|^2) dx \\ & + \frac{1}{\zeta} \int_{\Omega} \rho_n|v_n|^2 dx + \frac{4\delta\varepsilon}{\beta} \int_{\Omega} |\nabla \rho_n^{\beta/2}|^2 dx + \frac{4\varepsilon}{\gamma} \int_{\Omega} |\nabla \rho_n^{\gamma/2}|^2 dx = \int_{\Omega} \rho_n \nabla c_n \cdot v_n dx. \end{aligned} \tag{16}$$

We estimate the right-hand side by integrating by parts and using equation (12) for  $\rho_n$ :

$$\begin{aligned} \int_{\Omega} \rho_n \nabla c_n \cdot v_n dx &= - \int_{\Omega} c_n \operatorname{div}(\rho_n v_n) dx = \int_{\Omega} c_n (\partial_t \rho_n - \varepsilon \Delta \rho_n) dx \\ &= \frac{d}{dt} \int_{\Omega} \rho_n c_n dx - \int_{\Omega} \rho_n \partial_t c_n dx - \varepsilon \int_{\Omega} c_n \Delta \rho_n dx. \end{aligned} \tag{17}$$

Taking into account the second equation in (12), the second term on the right-hand side is written as

$$\begin{aligned} - \int_{\Omega} \rho_n \partial_t c_n dx &= - \int_{\Omega} (\partial_t c_n - \Delta c_n + c_n) \partial_t c_n dx \\ &= - \int_{\Omega} |\partial_t c_n|^2 dx - \frac{1}{2} \frac{d}{dt} \int_{\Omega} (|\nabla c_n|^2 + c_n^2) dx. \end{aligned}$$

Because of  $\rho_n c_n \geq 0$ , the last term on the right-hand side of (17) becomes

$$\begin{aligned} -\varepsilon \int_{\Omega} c_n \Delta \rho_n dx &= -\varepsilon \int_{\Omega} \rho_n \Delta c_n dx = -\varepsilon \int_{\Omega} \rho_n (\partial_t c_n + c_n - \rho_n) dx \\ &\leq -\varepsilon \int_{\Omega} \rho_n \partial_t c_n dx + \varepsilon \int_{\Omega} \rho_n^2 dx \leq \frac{\varepsilon}{4} \int_{\Omega} |\partial_t c_n|^2 dx + 2\varepsilon \int_{\Omega} \rho_n^2 dx \\ &\leq \frac{\varepsilon}{4} \int_{\Omega} |\partial_t c_n|^2 dx + 2\varepsilon \int_{\Omega} \rho_n^\gamma dx + C(\gamma, \Omega)\varepsilon, \end{aligned}$$

where the last inequality follows from  $\gamma \geq 2$ . (We observe that at this point, we can weaken the condition to  $\gamma > 8/5$  by using the Gagliardo–Nirenberg inequality and the estimate for  $\|\nabla \rho_n^{\gamma/2}\|_{L^2(\Omega)}$  from (16).) We insert these estimates into (17):

$$\begin{aligned} \int_{\Omega} \rho_n \nabla c_n \cdot v_n dx &\leq - \frac{d}{dt} \int_{\Omega} \left( \frac{1}{2} (|\nabla c_n|^2 + c_n^2) - \rho_n c_n \right) dx - \int_{\Omega} |\partial_t c_n|^2 dx \\ &+ \frac{\varepsilon}{4} \int_{\Omega} |\partial_t c_n|^2 dx + 2\varepsilon \int_{\Omega} \rho_n^\gamma dx + C(\gamma, \Omega)\varepsilon. \end{aligned}$$

Therefore, (16) leads to

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \left( \frac{1}{2}\rho_n|v_n|^2 + \psi(\rho_n) + \frac{1}{2} (|\nabla c_n|^2 + c_n^2) - \rho_n c_n + \frac{\delta}{\beta-1}\rho_n^\beta \right) dx \\ & + \int_{\Omega} (\mu|\nabla v_n|^2 + (\lambda + \mu)|\operatorname{div} v_n|^2) dx + \frac{4\delta\varepsilon}{\beta} \int_{\Omega} |\nabla \rho_n^{\beta/2}|^2 dx + \frac{4\varepsilon}{\gamma} \int_{\Omega} |\nabla \rho_n^{\gamma/2}|^2 dx \\ & + \left( 1 - \frac{\varepsilon}{4} \right) |\partial_t c_n|^2 dx + \frac{1}{\zeta} \int_{\Omega} \rho_n|v_n|^2 dx \leq 2\varepsilon \int_{\Omega} \rho_n^\gamma dx + C\varepsilon, \end{aligned} \tag{18}$$

where  $C > 0$  only depends on  $\gamma$  and  $\operatorname{meas}(\Omega)$  but is independent of  $n, \delta$ , and  $\varepsilon$ . This proves the inequality for  $E(\rho_n, v_n, c_n)$ .

*Step 2: Energy inequality for  $H$ .* We need to estimate  $\int_{\Omega} \rho_n c_n dx$  in  $E(\rho_n, v_n, c_n)$ . By Lemma 9 in Appendix 4, applied to  $m = \gamma$ ,  $\kappa = 1/(2(\gamma - 1))$ , and  $\xi = 1/4$ ,

$$\int_{\Omega} \rho_n c_n dx \leq \frac{1}{2(\gamma - 1)} \|\rho_n\|_{L^\gamma(\Omega)}^\gamma + \frac{1}{4} \|\nabla c_n\|_{L^2(\Omega)}^2 + C_1(\gamma) \|c_n\|_{L^1(\Omega)}^{C_2(\gamma)}. \tag{19}$$

Equation (12) implies that the mass is conserved,  $\|\rho_n(t)\|_{L^1(\Omega)} = \|\rho_\delta^0\|_{L^1(\Omega)}$  for  $0 < t < T_n$ . Furthermore, by the second equation in (12),

$$\frac{d}{dt} \int_{\Omega} c_n dx = \int_{\Omega} \rho_n dx - \int_{\Omega} c_n dx.$$

This is an ordinary differential equation for  $t \mapsto \|c_n(t)\|_{L^1(\Omega)}$ , and a comparison principle as well as the nonnegativity of  $c_n$  imply that

$$\|c_n(t)\|_{L^1(\Omega)} = \int_{\Omega} c_n dx \leq \max \left\{ \int_{\Omega} c^0 dx, \int_{\Omega} \rho_\delta^0 dx \right\} \leq C,$$

where  $C > 0$  is independent of  $\delta$ . Thus, we conclude from (19) and  $\rho_n^\gamma/(2(\gamma - 1)) = \frac{1}{2}\psi(\rho_n)$  that

$$\int_{\Omega} \rho_n c_n dx \leq \frac{1}{2} \int_{\Omega} \psi(\rho_n) dx + \frac{1}{4} \|\nabla c_n\|_{L^2(\Omega)}^2 + C(\rho^0, c^0).$$

It follows from the definitions of  $E(\rho_n, v_n, c_n)$  and  $H(\rho_n, v_n, c_n)$  that

$$\begin{aligned} E(\rho_n, v_n, c_n) &\geq \int_{\Omega} \left( \frac{1}{2} \psi(\rho_n) + \frac{1}{2} \rho_n |v_n|^2 + \frac{1}{4} |\nabla c_n|^2 + \frac{1}{2} c_n^2 \right) dx - C(\rho^0, c^0) \\ &= H(\rho_n, v_n, c_n) - C(\rho^0, c^0). \end{aligned}$$

We insert these estimates in (18) and integrate over  $(0, t)$  for  $0 < t < T_n$ :

$$\begin{aligned} &H((\rho_n, v_n, c_n)(t)) + \int_0^t \int_{\Omega} (\mu |\nabla v_n|^2 + (\lambda + \mu) |\operatorname{div} v_n|^2) dx ds + \frac{4\delta\varepsilon}{\beta} \int_0^t \int_{\Omega} |\nabla \rho_n^{\beta/2}|^2 dx ds \\ &+ \frac{4\varepsilon}{\gamma} \int_0^t \int_{\Omega} |\nabla \rho_n^{\gamma/2}|^2 dx ds + \left(1 - \frac{\varepsilon}{4}\right) \int_0^t \int_{\Omega} |\partial_s c_n|^2 dx ds + \frac{1}{\zeta} \int_0^t \int_{\Omega} \rho_n |v_n|^2 dx ds \\ &\leq E(\rho_\delta^0, v^0, c^0) + C\varepsilon \int_0^t \int_{\Omega} H(\rho_n, v_n, c_n) dx ds + C(\rho^0, c^0) + C\varepsilon t, \end{aligned}$$

where we used  $\int_{\Omega} \rho_n^\gamma dx \leq CH(\rho_n, v_n, c_n)$ . An application of Gronwall’s lemma finishes the proof.  $\square$

Lemma 3 allows us to conclude as in [8, Section 2.3] that  $T = T_n$ . Moreover, it yields the following estimates uniform in  $(n, \delta, \varepsilon)$ :

$$\begin{aligned} (\rho_n) &\text{ is uniformly bounded in } L^\infty(0, T; L^\gamma(\Omega)), \\ (\sqrt{\rho_n} v_n) &\text{ is uniformly bounded in } L^\infty(0, T; L^2(\Omega; \mathbb{R}^3)), \\ (\nabla v_n) &\text{ is uniformly bounded in } L^2(0, T; L^2(\Omega; \mathbb{R}^{3 \times 3})), \\ (c_n) &\text{ is uniformly bounded in } L^\infty(0, T; H^1(\Omega)) \cap H^1(0, T; L^2(\Omega)). \end{aligned} \tag{20}$$

**2.3. Limit  $(n, \delta, \varepsilon) \rightarrow (\infty, 0, 0)$**

The limit  $n \rightarrow \infty$  can be performed as in [1, Section 2.3] via the Aubin–Lions compactness lemma. This gives a solution  $(\rho_\delta, v_\delta, c_\delta)$  to (12)–(15). It satisfies the energy inequalities in Lemma 3. In particular, we conclude a uniform bound for  $\rho_\delta$  in  $L^\infty(0, T; L^\gamma(\Omega))$ . By the existence theory for the compressible Navier–Stokes equations, we can pass to the limit  $(\delta, \varepsilon) \rightarrow 0$ ; see, e.g., [5, 16]. Indeed, to derive a uniform estimate for the mass density in  $L^{\gamma+\theta}(\Omega \times (0, T))$  for some  $\theta > 0$ , we need to use the test function

$$\phi_B = \mathcal{B} \left( \rho_\delta^\theta - \frac{1}{|\Omega|} \int_{\Omega} \rho_\delta^\theta dx \right),$$

in the weak formulation of the approximate momentum equation, where  $\mathcal{B}$  is the Bogovskii operator [16, Section 3.3.1.2]. Compared to the compressible Navier–Stokes equations, the momentum equation includes the chemotaxis term  $\rho_\delta \nabla c_\delta$ , which needs to be estimated. This means that we need a bound for

$$I = \int_0^T \int_\Omega \rho_\delta \nabla c_\delta \cdot \phi_B dx dt. \tag{21}$$

Using the second equation in (12),

$$\rho_\delta \nabla c_\delta = (\partial_t c_\delta - \Delta c_\delta + c_\delta) \nabla c_\delta = (\partial_t c_\delta + c_\delta) \nabla c_\delta - \operatorname{div}(\nabla c_\delta \otimes \nabla c_\delta) + \frac{1}{2} \nabla |\nabla c_\delta|^2,$$

we can write  $I = I_1 + \dots + I_4$ , where

$$\begin{aligned} I_1 &= \int_0^T \int_\Omega \partial_t c_\delta \nabla c_\delta \cdot \phi_B dx dt, & I_2 &= \int_0^T \int_\Omega c_\delta \nabla c_\delta \cdot \phi_B dx dt, \\ I_3 &= \int_0^T \int_\Omega \nabla c_\delta \otimes \nabla c_\delta : \nabla \phi_B dx dt, & I_4 &= -\frac{1}{2} \int_0^T \int_\Omega |\nabla c_\delta|^2 \operatorname{div} \phi_B dx dt. \end{aligned}$$

We start with the term  $I_1$ . First, let  $\gamma > 2$ . By parabolic regularity theory (see Theorem 10 in the Appendix with  $p = q = \gamma$ ), the continuous embedding  $W^{2-2/\gamma, \gamma}(\Omega) \hookrightarrow W^{1, \gamma}(\Omega)$  and the second equation in (12) yield

$$\begin{aligned} \|\nabla c_\delta\|_{L^2(0, T; L^\gamma(\Omega))} &\leq C \|c_\delta\|_{L^2(0, T; W^{2-2/\gamma, \gamma}(\Omega))} \\ &\leq C (\|\rho_\delta\|_{L^\infty(0, T; L^\gamma(\Omega))} + \|c^0\|_{W^{2-2/\gamma, \gamma}(\Omega)}) \leq C. \end{aligned} \tag{22}$$

Hence, using Hölder’s inequality, the assumption  $\gamma > 2$ , and the previous estimates as well as the uniform estimates from the energy inequality,

$$I_1 \leq \|\partial_t c_\delta\|_{L^2(0, T; L^2(\Omega))} \|\nabla c_\delta\|_{L^2(0, T; L^\gamma(\Omega))} \|\phi_B\|_{L^\infty(0, T; L^{2\gamma/(\gamma-2)}(\Omega))} \leq C \|\phi_B\|_{L^\infty(0, T; W^{1, r}(\Omega))},$$

where  $r = 6\gamma/(5\gamma - 6)$  is such that  $W^{1, r}(\Omega) \hookrightarrow L^{2\gamma/(\gamma-2)}(\Omega)$ . We deduce from the boundedness of  $\mathcal{B} : L_0^r(\Omega) \rightarrow W_0^{1, r}(\Omega)$  for  $1 < r < \infty$ , where  $L_0^r(\Omega)$  is the space of all  $L^r(\Omega)$  functions  $u$  satisfying  $\int_\Omega u dx = 0$ , that

$$\begin{aligned} I_1 &\leq C \left\| \rho_\delta^\theta - \frac{1}{|\Omega|} \int_\Omega \rho_\delta^\theta dx \right\|_{L^\infty(0, T; L^r(\Omega))} \leq C \|\rho_\delta^\theta\|_{L^\infty(0, T; L^r(\Omega))} \\ &\leq C \|\rho_\delta\|_{L^\infty(0, T; L^{r\theta}(\Omega))}^\theta \leq C \|\rho_\delta\|_{L^\infty(0, T; L^\gamma(\Omega))}^\theta \leq C. \end{aligned}$$

The last but one step follows if  $r\theta \leq \gamma$ , which requires the choice  $0 < \theta \leq 5\gamma/6 - 1$ , and the last step is a consequence of the energy inequality.

Next, let  $3/2 < \gamma \leq 2$ . We apply Theorem 10 with  $p = 2, q = \gamma$  to find that

$$\|c_\delta\|_{L^2(0, T; W^{2, \gamma}(\Omega))} + \|\partial_t c_\delta\|_{L^2(0, T; L^\gamma(\Omega))} \leq C (\|\rho_\delta\|_{L^2(0, T; L^\gamma(\Omega))} + \|c^0\|_{W^{1, \gamma}(\Omega)}) \leq C.$$

Hence, we deduce from the continuous embedding  $W^{2, \gamma}(\Omega) \hookrightarrow W^{1, 3\gamma/(3-\gamma)}(\Omega)$  that

$$\begin{aligned} I_1 &\leq \|\partial_t c_\delta\|_{L^2(0, T; L^\gamma(\Omega))} \|\nabla c_\delta\|_{L^2(0, T; L^{3\gamma/(3-\gamma)}(\Omega))} \|\phi_B\|_{L^\infty(0, T; L^{3\gamma/(4\gamma-6)}(\Omega))} \\ &\leq C \|\partial_t c_\delta\|_{L^2(0, T; L^\gamma(\Omega))} \|c_\delta\|_{L^2(0, T; W^{2, \gamma}(\Omega))} \|\phi_B\|_{L^\infty(0, T; W^{1, r}(\Omega))}, \end{aligned}$$

where now  $r = 3\gamma/(5\gamma - 6)$ . We choose  $\theta > 0$  such that  $r\theta \leq \gamma$ , which is equivalent to  $\theta \leq 5\gamma/3 - 2$ , and we can choose  $\theta > 0$  satisfying this inequality. Then, arguing as in the case  $\gamma \geq 2$ ,

$$I_1 \leq C \|\rho_\delta\|_{L^\infty(0, T; L^{r\theta}(\Omega))}^\theta \leq C \|\rho_\delta\|_{L^\infty(0, T; L^\gamma(\Omega))}^\theta \leq C.$$

For the term  $I_3$ , we consider again first the case  $\gamma > 2$ :

$$I_3 \leq \|\nabla c_\delta\|_{L^2(0, T; L^\gamma(\Omega))}^2 \|\nabla \phi_B\|_{L^\infty(0, T; L^r(\Omega))} \leq C \|\rho_\delta\|_{L^\infty(0, T; L^{r\theta}(\Omega))}^\theta \leq C,$$



where  $r = \gamma/(\gamma - 2)$ , and the last inequality follows if  $r\theta \leq \gamma$ , which is equivalent to  $\theta \leq \gamma - 2$ . If  $3/2 < \gamma \leq 2$ , we proceed similarly as for  $I_1$ :

$$I_3 \leq \|\nabla c_\delta\|_{L^2(0,T;L^{3\gamma/(3-\gamma)}(\Omega))}^2 \|\nabla \phi_B\|_{L^\infty(0,T;L^r(\Omega))} \leq C \|\rho_\delta\|_{L^\infty(0,T;L^{r\theta}(\Omega))}^\theta \leq C,$$

where  $r = \gamma/(2\gamma - 3)$  and we need  $r\theta \leq \gamma$  or, equivalently,  $\theta \leq 2\gamma - 3$ .

The term  $I_2$  is estimated in a similar way as  $I_1$ , and  $I_4$  can be bounded as  $I_3$ . This shows that  $I$  is bounded and provides a uniform estimate for  $\rho_\delta$  in  $L^{\gamma+\theta}(\Omega \times (0, T))$ . Now we can proceed as in [16, Section 7.3] to prove the strong convergence of the pressure.

*Remark 4.* (On the condition on  $\gamma$  in [1]) Aïssa and Alexandre have estimated the term  $I$ , defined in (21), in a different way. They used the test function  $\psi_B = \mathcal{B}(\rho_\delta - |\Omega|^{-1} \int_\Omega \rho_\delta dx)$ :

$$\begin{aligned} I &\leq \|\rho_\delta\|_{L^2(\Omega \times (0, T))} \|\nabla c_\delta\|_{L^2(\Omega \times (0, T))} \|\psi_B\|_{L^\infty(0, T; L^\infty(\Omega))} \\ &\leq \|\rho_\delta\|_{L^2(\Omega \times (0, T))} \|\nabla c_\delta\|_{L^2(\Omega \times (0, T))} \|\rho_\delta\|_{L^\infty(0, T; L^r(\Omega))}, \end{aligned}$$

which is a consequence of the estimate  $\|\mathcal{B}(f)\|_{L^\infty(\Omega)} \leq C \|\mathcal{B}(f)\|_{W^{1,r}(\Omega)} \leq C \|f\|_{L^r(\Omega)}$  choosing  $r > d = 3$ . Thus, the technique of [1] only works for  $\gamma > 3$ . □

*Remark 5.* (On the condition  $\gamma > 8/5$ ) This restriction is needed to estimate the integral  $\int_\Omega \rho_n c_n dx$  by means of Lemma 9. The idea is to obtain “small” terms that can be absorbed by the left-hand side of the energy inequality (18) and terms that can be controlled (the  $L^1(\Omega)$  norm of  $c_n$ ). By the Hölder and Gagliardo–Nirenberg inequalities, we may estimate in a different way:

$$\int_\Omega \rho_n c_n dx \leq \|\rho_n\|_{L^\gamma(\Omega)} \|c_n\|_{L^{\gamma/(\gamma-1)}(\Omega)} \leq C \|\rho_n\|_{L^\gamma(\Omega)} \|c_n\|_{W^{2,\gamma}(\Omega)}^\theta \|c_n\|_{L^1(\Omega)}^{1-\theta},$$

where  $\theta = 3/(5\gamma - 3) \in (0, 1)$  (which requires that  $\gamma > 6/5$ ). It follows from the maximal regularity result of Theorem 10 that

$$\int_\Omega \rho_n c_n dx \leq C \|\rho_n\|_{L^\gamma(\Omega)} (\|\rho_n\|_{L^\gamma(\Omega)} + 1)^\theta,$$

where  $C > 0$  depends on  $\|c^0\|_{L^1(\Omega)}$ . We can conclude if  $1 + \theta < \gamma$ , which is equivalent to  $\gamma > 8/5$ . Thus, even taking into account maximal regularity does not improve the range for  $\gamma$ .

*Remark 6.* (Improving the condition on  $\gamma$ ) When the dynamics of the chemical concentration is much faster than that one of the cell density, we can neglect the time derivative of the concentration in (3), and  $c_\delta$  solves  $0 = \Delta c_\delta - c_\delta + \rho_\delta$  in  $\Omega$ . In this situation, we are able to weaken the condition on  $\gamma$  to  $\gamma > 3/2$ . Indeed, estimate (22) still holds for the elliptic problem. The embedding  $W^{1,\gamma}(\Omega) \hookrightarrow L^{3\gamma/(3-\gamma)}(\Omega)$  for  $\gamma < 3$  then shows that

$$\|\nabla c_\delta\|_{L^\infty(0,T;L^{3\gamma/(3-\gamma)}(\Omega))} \leq C \|c_\delta\|_{L^\infty(0,T;W^{2,\gamma}(\Omega))} \leq C \|\rho_\delta\|_{L^\infty(0,T;L^\gamma(\Omega))} \leq C.$$

Hence, using Hölder’s inequality, we estimate

$$\begin{aligned} I_3 &\leq \|\nabla c_\delta\|_{L^\infty(0,T;L^{3\gamma/(3-\gamma)}(\Omega))}^2 \|\nabla \phi_B\|_{L^\infty(0,T;L^{3\gamma/(5\gamma-6)}(\Omega))} \\ &\leq C \|\rho_\delta\|_{L^\infty(0,T;L^\gamma(\Omega))}^2 \|\rho_\delta\|_{L^\infty(0,T;L^{3\gamma/(5\gamma-6)}(\Omega))}^\theta \leq C + C \|\rho_\delta\|_{L^\infty(0,T;L^\gamma(\Omega))}^{2+\theta} \leq C, \end{aligned}$$

provided that  $0 < \theta < (5\gamma - 6)/3$ . The terms  $I_2$  and  $I_4$  are estimated in a similar way and  $I_1 = 0$ , thus proving that  $I$  is bounded. This yields a uniform estimate for  $\rho_\delta$  in  $L^{\gamma+\theta}(\Omega \times (0, T))$  for  $\gamma > 3/2$  according to the theory of the compressible Navier–Stokes equations. □

### 3. Relative Energy Inequality

We show a relative energy inequality for smooth functions.

**Lemma 7** (Relative energy inequality). *Let  $(\rho, v, c)$  be a smooth solution to (1)–(5) and let  $(r, u, z)$  be smooth functions satisfying  $r > 0$  in  $\bar{\Omega} \times [0, T]$  and  $u = 0$  on  $\partial\Omega$ . Then the relative energy inequality (10) holds for  $0 < t < T$  with*

$$\begin{aligned} R(\rho, v, c|r, u, z) &= - \int_{\Omega} p(\rho|r) \operatorname{div} u dx - \int_{\Omega} \psi''(r)(\rho - r)g dx - \int_{\Omega} h\partial_t(c - z)dx \\ &\quad - \int_{\Omega} \nabla(c - z) \cdot ((\rho - r)u)dx + \int_{\Omega} (c - z)g dx \\ &\quad - \int_{\Omega} \rho(v - u) \otimes (v - u) : \nabla u dx - \frac{1}{\zeta} \int_{\Omega} \rho|v - u|^2 dx \\ &\quad - \int_{\Omega} \left( \frac{\rho - r}{r} (\mu\Delta u + (\lambda + \mu)\nabla \operatorname{div} u) + \rho f \right) \cdot (v - u) dx, \end{aligned}$$

where

$$f = \partial_t u + u \cdot \nabla u + \frac{1}{r} \nabla p(r) - \frac{1}{r} (\mu\Delta u + (\lambda + \mu)\nabla \operatorname{div} u) - \nabla z + \frac{u}{\zeta}, \tag{23}$$

$$g = \partial_t r + \operatorname{div}(ru), \quad h = \partial_t z - \Delta z + z - r. \tag{24}$$

We prove in Sect. 4 that the relative energy inequality (10) holds for finite energy weak solutions  $(\rho, v, c)$  and  $(\bar{\rho}, \bar{v}, \bar{c})$ , where  $(\bar{\rho}, \bar{v})$  satisfies (9). The proof of (10) follows the lines of [9, Section 3.2], but some steps are different due to the additional chemotaxis force. For this reason, and for the convenience of the reader, we present a full proof.

*Proof.* Let  $(r_m, u_m, z_m)_{m \in \mathbb{N}}$  be smooth functions satisfying  $r_m > 0$  in  $\bar{\Omega} \times [0, T]$ ,  $v_m \in C^1([0, T]; X_m)$ , and  $v_m = 0$  on  $\partial\Omega$  such that  $(r_m, u_m, z_m) \rightarrow (r, z, u)$  as  $m \rightarrow \infty$  in a sense made precise in Step 3 below. Here,  $X_m$  is the Faedo–Galerkin space defined in Sect. 2.1. We introduce

$$\begin{aligned} f_m &= \partial_t u_m + u_m \cdot \nabla u_m + \frac{1}{r_m} \nabla p(r_m) - \frac{1}{r_m} (\mu\Delta u_m + (\lambda + \mu)\nabla \operatorname{div} u_m) \\ &\quad - \nabla z_m + \frac{u_m}{\zeta}, \end{aligned} \tag{25}$$

$$g_m = \partial_t r_m + \operatorname{div}(r_m u_m), \quad h_m = \partial_t z_m - \Delta z_m + z_m - r_m. \tag{26}$$

Then  $(f_m, g_m, h_m) \rightarrow (f, g, h)$  as  $m \rightarrow \infty$  in the sense of distributions, where  $(f, g, h)$  is defined in (23)–(24). Finally, let  $(\rho_n, v_n, c_n)$  be a Galerkin solution to (12)–(15). We compute in the following the approximate relative energy inequality.

*Step 1: Time derivative of the relative kinetic energy.* We derive an equation for the time evolution of the relative kinetic energy  $\frac{1}{2} \int_{\Omega} \rho_n |v_n - u_m|^2 dx$ . It follows from the approximative mass balance equation (12) that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} (\rho_n |v_n - u_m|^2) &= -\frac{1}{2} (\operatorname{div}(\rho_n v_n) - \varepsilon \Delta \rho_n) |v_n - u_m|^2 + \rho_n \partial_t (v_n - u_m) \cdot (v_n - u_m) \\ &= -\frac{1}{2} \operatorname{div}(\rho_n v_n |v_n - u_m|^2) + \rho_n v_n \cdot \nabla (v_n - u_m) \cdot (v_n - u_m) \\ &\quad + \rho_n \partial_t (v_n - u_m) \cdot (v_n - u_m) + \frac{\varepsilon}{2} \Delta \rho_n |v_n - u_m|^2. \end{aligned} \tag{27}$$

Since  $\rho_n (\partial_t v_n + v_n \cdot \nabla v_n) = \partial_t (\rho_n v_n) + \operatorname{div}(\rho_n v_n \otimes v_n) - \varepsilon \Delta \rho_n v_n$ , the second and third terms on the right-hand side are written as

$$\begin{aligned} &\rho_n v_n \cdot \nabla (v_n - u_m) \cdot (v_n - u_m) + \rho_n \partial_t (v_n - u_m) \cdot (v_n - u_m) \\ &= \rho_n (\partial_t v_n + v_n \cdot \nabla v_n) \cdot (v_n - u_m) - \rho_n (\partial_t u_m + v_n \cdot \nabla u_m) \cdot (v_n - u_m) \\ &= (\partial_t (\rho_n v_n) + \operatorname{div}(\rho_n v_n \otimes v_n)) \cdot (v_n - u_m) - \varepsilon \Delta \rho_n v_n \cdot (v_n - u_m) \\ &\quad - \rho_n (\partial_t u_m + u_m \cdot \nabla u_m) \cdot (v_n - u_m) - \rho_n (v_n - u_m) \cdot \nabla u_m \cdot (v_n - u_m). \end{aligned}$$

We insert this expression into (27), integrate over  $\Omega$ , and replace  $\partial_t(\rho_n v_n) + \operatorname{div}(\rho_n v_n \otimes v_n)$  by the momentum equation (13):

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho_n |v_n - u_m|^2 dx &= \frac{\varepsilon}{2} \int_{\Omega} \Delta \rho_n |v_n - u_m|^2 dx - \int_{\Omega} \nabla(p(\rho_n) + \delta \rho_n^\beta) \cdot (v_n - u_m) dx \\ &\quad - \int_{\Omega} (\mu \nabla v_n \cdot \nabla(v_n - u_m) + (\lambda + \mu) \operatorname{div} v_n \operatorname{div}(v_n - u_m)) dx \\ &\quad - \varepsilon \int_{\Omega} \nabla \rho_n \cdot \nabla v_n \cdot (v_n - u_m) dx + \int_{\Omega} \rho_n \nabla c_n \cdot (v_n - u_m) dx \\ &\quad - \frac{1}{\zeta} \int_{\Omega} \rho_n v_n \cdot (v_n - u_m) dx - \varepsilon \int_{\Omega} \Delta \rho_n v_n \cdot (v_n - u_m) dx \\ &\quad - \int_{\Omega} \rho_n (\partial_t u_m + u_m \cdot \nabla u_m) \cdot (v_n - u_m) dx - \int_{\Omega} \rho_n (v_n - u_m) \cdot \nabla u_m \cdot (v_n - u_m) dx. \end{aligned} \tag{28}$$

We wish to reformulate the last but one term in the previous equality. For this, we add and subtract  $r_m$  and replace  $r_m(\partial_t u_m + u_m \cdot \nabla u_m)$  by (25):

$$\begin{aligned} & - \int_{\Omega} \rho_n (\partial_t u_m + u_m \cdot \nabla u_m) \cdot (v_n - u_m) dx \\ &= - \int_{\Omega} \left(1 + \frac{\rho_n - r_m}{r_m}\right) (r_m(\partial_t u_m + u_m \cdot \nabla u_m)) \cdot (v_n - u_m) dx \\ &= \int_{\Omega} \left(1 + \frac{\rho_n - r_m}{r_m}\right) \left(\nabla p(r_m) - r_m \nabla z_m + \frac{r_m u_m}{\zeta} - r_m f_m\right) \cdot (v_n - u_m) dx \\ &\quad - \int_{\Omega} \left(1 + \frac{\rho_n - r_m}{r_m}\right) (\mu \Delta u_m + (\lambda + \mu) \nabla \operatorname{div} u_m) \cdot (v_n - u_m) dx. \end{aligned}$$

Then, after a computation, (28) becomes

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho_n |v_n - u_m|^2 dx &= - \int_{\Omega} \left(\nabla p(\rho_n) - \frac{\rho_n}{r_m} \nabla p(r_m)\right) \cdot (v_n - u_m) dx \\ &\quad - \delta \int_{\Omega} \nabla \rho_n^\beta \cdot (v_n - u_m) dx + \varepsilon \int_{\Omega} \nabla \rho_n \cdot \nabla u_m \cdot (v_n - u_m) dx \\ &\quad - \int_{\Omega} (\mu |\nabla(v_n - u_m)|^2 + (\lambda + \mu) |\operatorname{div}(v_n - u_m)|^2) dx \\ &\quad - \int_{\Omega} \rho_n (v_n - u_m) \otimes (v_n - u_m) : \nabla u_m dx + \int_{\Omega} \rho_n \nabla(c_n - z_m) \cdot (v_n - u_m) dx \\ &\quad - \frac{1}{\zeta} \int_{\Omega} \rho_n |v_n - u_m|^2 dx - \int_{\Omega} \rho_n f_m \cdot (v_n - u_m) dx \\ &\quad - \int_{\Omega} \frac{\rho_n - r_m}{r_m} (\mu \Delta u_m + (\lambda + \mu) \nabla \operatorname{div} u_m) \cdot (v_n - u_m) dx. \end{aligned} \tag{29}$$

We rewrite the first, second, and sixth terms on the right-hand side of (28).

*Step 2a: Reformulation of the pressure term.* Observing that  $p'(z) = z\psi''(z)$  for  $z \geq 0$  (see (11)) and that  $\rho_m - r_m$  satisfies

$$\partial_t(\rho_n - r_m) + \operatorname{div}((\rho_n - u_m)u_m + \rho_n(v_n - u_m)) = \varepsilon \Delta \rho_n - g_m,$$

we can write the first term on the right-hand side of (29) as

$$\begin{aligned} - \int_{\Omega} \left(\nabla p(\rho_n) - \frac{\rho_n}{r_m} \nabla p(r_m)\right) \cdot (v_n - u_m) dx &= \int_{\Omega} \rho_n \nabla(\psi'(\rho_n) - \psi'(r_m)) \cdot (v_n - u_m) dx \\ &= - \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \operatorname{div}(\rho_n(v_n - u_m)) dx \end{aligned}$$

$$\begin{aligned}
 &= - \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \partial_t(\rho_n - r_m) dx \\
 &\quad - \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \operatorname{div}((\rho_n - r_m)u_m) dx \\
 &\quad + \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) (\varepsilon \Delta \rho_n - g_m) dx.
 \end{aligned} \tag{30}$$

Taking into account that the evolution of the relative internal energy is given by

$$\begin{aligned}
 \partial_t \psi(\rho_n | r_m) &= \partial_t (\psi(\rho_n) - \psi(r_m) - \psi'(r_m)(\rho_n - r_m)) \\
 &= (\psi'(\rho_n) - \psi'(r_m)) \partial_t \rho_n - \psi''(r_m) \partial_t r_m (\rho_n - r_m),
 \end{aligned}$$

the first term on the right-hand side of (30) is reformulated as

$$\begin{aligned}
 &- \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \partial_t(\rho_n - r_m) dx \\
 &= - \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \partial_t \rho_n dx + \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \partial_t r_m dx \\
 &= - \int_{\Omega} \left( \frac{d}{dt} \psi(\rho_n | r_m) + \psi''(r_m) \partial_t r_m (\rho_n - r_m) \right) dx + \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \partial_t r_m dx \\
 &= - \frac{d}{dt} \int_{\Omega} \psi(\rho_n | r_m) dx + \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m) - \psi''(r_m)(\rho_n - r_m)) (g_m - \operatorname{div}(r_m u_m)) dx,
 \end{aligned}$$

where we used definition (26) of  $g_m$  in the last step. Integrating by parts to get rid of the divergence, inserting the corresponding expression into (30), and observing that the integral over  $(\psi'(\rho_n) - \psi'(r_m))g_m$  cancels with the corresponding expression in (30), we find that

$$\begin{aligned}
 &- \int_{\Omega} \left( \nabla p(\rho_n) - \frac{\rho_n}{r_m} \nabla p(r_m) \right) \cdot (v_n - u_m) dx = - \frac{d}{dt} \int_{\Omega} \psi(\rho_n | r_m) dx \\
 &\quad + \int_{\Omega} \{ \nabla (\psi'(\rho_n) - \psi'(r_m)) \cdot (\rho_n u_m) - \nabla (\psi''(r_m)(\rho_n - r_m)) \cdot (r_m u_m) \} dx \\
 &\quad + \varepsilon \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \Delta \rho_n dx - \int_{\Omega} \psi''(r_m)(\rho_n - r_m) g_m dx.
 \end{aligned}$$

We claim that the second term on the right-hand side can be formulated in terms of the relative pressure  $p(\rho_n | r_m) = p(\rho_n) - p(r_m) - p'(r_m)(\rho_n - r_m)$ . It follows from (11) that  $\nabla p(\rho_n) = \rho_n \nabla \psi'(\rho_n)$ ,  $\nabla p'(r_m) = \nabla \psi'(r_m) + r_m \nabla \psi''(r_m)$  and hence,

$$\begin{aligned}
 \nabla p(\rho_n | r_m) &= \rho_n \nabla \psi'(\rho_n) - r_m \nabla \psi'(r_m) - \nabla (r_m \psi''(r_m)(\rho_n - r_m)) \\
 &= \rho_n \nabla (\psi'(\rho_n) - \psi'(r_m)) - r_m \nabla (\psi''(r_m)(\rho_n - r_m))
 \end{aligned}$$

and consequently,

$$\begin{aligned}
 &\int_{\Omega} \{ \nabla (\psi'(\rho_n) - \psi'(r_m)) \cdot (\rho_n u_m) - \nabla (\psi''(r_m)(\rho_n - r_m)) \} \cdot (r_m u_m) dx \\
 &= \int_{\Omega} \nabla p(\rho_n | r_m) \cdot u_m dx = - \int_{\Omega} p(\rho_n | r_m) \operatorname{div} u_m dx.
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 &- \int_{\Omega} \left( \nabla p(\rho_n) - \frac{\rho_n}{r_m} \nabla p(r_m) \right) \cdot (v_n - u_m) dx = - \frac{d}{dt} \int_{\Omega} \psi(\rho_n | r_m) dx \\
 &\quad - \int_{\Omega} p(\rho_n | r_m) \operatorname{div} u_m dx + \varepsilon \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \Delta \rho_n dx \\
 &\quad - \int_{\Omega} \psi''(r_m)(\rho_n - r_m) g_m dx.
 \end{aligned} \tag{31}$$

*Step 2b: Reformulation of the chemotaxis term.* We reformulate the sixth term on the right-hand side of (29) by integrating by parts and using the mass balances (12) and (26):

$$\begin{aligned}
 \int_{\Omega} \rho_n \nabla(c_n - z_m) \cdot (v_n - u_m) dx &= - \int_{\Omega} (c_n - z_m) \operatorname{div}(\rho_n(v_n - u_m)) dx \\
 &= \int_{\Omega} (c_n - z_m) \operatorname{div}(-\rho_n v_n + (\rho_n - r_m)u_m + r_m u_m) dx \\
 &= \int_{\Omega} (c_n - z_m) (\partial_t(\rho_n - r_m) + \operatorname{div}((\rho_n - r_m)u_m) - \varepsilon \Delta \rho_n + g_m) dx \\
 &= \frac{d}{dt} \int_{\Omega} (c_n - z_m)(\rho_n - r_m) dx - \int_{\Omega} (\rho_n - r_m) \partial_t(c_n - z_m) dx \\
 &\quad - \int_{\Omega} \nabla(c_n - z_m) \cdot ((\rho_n - r_m)u_m) dx - \int_{\Omega} (c_n - z_m)(\varepsilon \Delta \rho_n - g_m) dx.
 \end{aligned} \tag{32}$$

In view of the second equation in (26), we have

$$\rho_n - r_m = \partial_t(c_n - z_m) - \Delta(c_n - z_m) + (c_n - z_m) + h_m.$$

We insert this expression into the second term on the right-hand side of (32):

$$\begin{aligned}
 \int_{\Omega} \rho_n \nabla(c_n - z_m) \cdot (v_n - u_m) dx &= \frac{d}{dt} \int_{\Omega} (c_n - z_m)(\rho_n - r_m) dx - \int_{\Omega} |\partial_t(c_n - z_m)|^2 dx \\
 &\quad - \frac{1}{2} \frac{d}{dt} \int_{\Omega} (|\nabla(c_n - z_m)|^2 + (c_n - z_m)^2) dx - \int_{\Omega} h_m \partial_t(c_n - z_m) dx \\
 &\quad - \int_{\Omega} \nabla(c_n - z_m) \cdot ((\rho_n - r_m)u_m) dx - \int_{\Omega} (c_n - z_m)(\varepsilon \Delta \rho_n - g_m) dx.
 \end{aligned} \tag{33}$$

*Step 2c: Reformulation of the artificial pressure term.* We rewrite the second term on the right-hand side of (29) by integrating by parts and using the mass balance equation (12):

$$\begin{aligned}
 -\delta \int_{\Omega} \nabla \rho_n^\beta \cdot (v_n - u_m) dx &= -\beta \delta \int_{\Omega} \rho_n^{\beta-2} \nabla \rho_n \cdot (\rho_n v_n) dx - \delta \int_{\Omega} \rho_n^\beta \operatorname{div} u_m dx \\
 &= \frac{\beta \delta}{\beta - 1} \int_{\Omega} \rho_n^{\beta-1} \operatorname{div}(\rho_n v_n) dx - \delta \int_{\Omega} \rho_n^\beta \operatorname{div} u_m dx \\
 &= \frac{\beta \delta}{\beta - 1} \int_{\Omega} \rho_n^{\beta-1} (\varepsilon \Delta \rho_n - \partial_t \rho_n) dx - \delta \int_{\Omega} \rho_n^\beta \operatorname{div} u_m dx \\
 &= -\frac{\delta}{\beta - 1} \frac{d}{dt} \int_{\Omega} \rho_n^\beta dx - \beta \delta \int_{\Omega} \rho_n^{\beta-2} |\nabla \rho_n|^2 dx - \delta \int_{\Omega} \rho_n^\beta \operatorname{div} u_m dx.
 \end{aligned} \tag{34}$$

*Step 2d: Collecting the reformulations.* We include the reformulations (31), (33), and (34) into (29) to find that

$$\begin{aligned}
 \frac{d}{dt} \int_{\Omega} &\left\{ \frac{1}{2} \rho_n |v_n - u_m|^2 + \psi(\rho_n |r_m) + \frac{1}{2} (|\nabla(c_n - z_m)|^2 + (c_n - z_m)^2) \right. \\
 &\quad \left. - (c_n - z_m)(\rho_n - r_m) + \frac{\delta}{\beta - 1} \rho_n^\beta \right\} dx + \int_{\Omega} |\partial_t(c_n - z_m)|^2 dx \\
 &+ \beta \delta \int_{\Omega} \rho_n^{\beta-2} |\nabla \rho_n|^2 dx + \int_{\Omega} (\mu |\nabla(v_n - u_m)|^2 + (\lambda + \mu) |\operatorname{div}(v_n - u_m)|^2) dx \\
 &= - \int_{\Omega} p(\rho_n |r_m) \operatorname{div} u_m dx + \varepsilon \int_{\Omega} (\psi'(\rho_n) - \psi'(r_m)) \Delta \rho_n dx \\
 &\quad - \int_{\Omega} \psi''(r_m) (\rho_n - r_m) g_m dx - \int_{\Omega} h_m \partial_t(c_n - z_m) dx - \delta \int_{\Omega} \rho_n^\beta \operatorname{div} u_m dx \\
 &\quad - \int_{\Omega} \nabla(c_n - z_m) \cdot ((\rho_n - r_m)u_m) dx - \int_{\Omega} (c_n - z_m)(\varepsilon \Delta \rho_n - g_m) dx
 \end{aligned}$$

$$\begin{aligned}
 & + \varepsilon \int_{\Omega} \nabla \rho_n \cdot \nabla u_m \cdot (v_n - u_m) dx - \int_{\Omega} \rho_n (v_n - u_m) \otimes (v_n - u_m) : \nabla u_m dx \\
 & - \frac{1}{\zeta} \int_{\Omega} \rho_n |v_n - u_m|^2 dx - \int_{\Omega} \rho_n f_m \cdot (v_n - u_m) dx \\
 & - \int_{\Omega} \frac{\rho_n - r_m}{r_m} (\mu \Delta u_m + (\lambda + \mu) \nabla \operatorname{div} u_m) \cdot (v_n - u_m) dx.
 \end{aligned} \tag{35}$$

*Step 3: Limit  $(n, m) \rightarrow \infty$  and  $(\delta, \varepsilon) \rightarrow 0$ .* As mentioned in [9, Section 3.3], the limit in the approximate relative energy inequality (35) follows step by step the existence proof in [5, Chapter 7] or [16, Chapter 7]. In particular, we perform first the limit  $n \rightarrow \infty$  in the Faedo–Galerkin approximation  $(\rho_n, v_n, c_n) \rightarrow (\rho_{\varepsilon, \delta}, v_{\varepsilon, \delta}, c_{\varepsilon, \delta})$ . Then the functions  $(r_m, u_m, z_m)$  are replaced by smooth functions  $(r, u, z)$  using a density argument. Third, we pass to the limit  $(\rho_{\varepsilon, \delta}, v_{\varepsilon, \delta}, c_{\varepsilon, \delta}) \rightarrow (\rho_{\delta}, v_{\delta}, c_{\delta})$  as  $\varepsilon \rightarrow 0$  and  $(\rho_{\delta}, v_{\delta}, c_{\delta}) \rightarrow (\rho, v, c)$  as  $\delta \rightarrow 0$ .

In view of the bounds (20), we can pass to the limit  $n \rightarrow \infty$  and  $(\delta, \varepsilon) \rightarrow 0$  in (35). We assume that  $(r_m, u_m, z_m)$  converges to  $(r, u, z)$  as  $m \rightarrow \infty$  in such a way that the limit  $m \rightarrow \infty$  in (35) is possible. Then some integrals in (35) disappear and we end up with

$$\begin{aligned}
 & \frac{d}{dt} \int_{\Omega} \left( \psi(\rho|r) + \frac{1}{2} \rho |v - u|^2 + \frac{1}{2} (|\nabla(c - z)|^2 + (c - z)^2) - (\rho - r)(c - z) \right) dx \\
 & + \int_{\Omega} (\mu |\nabla(v - u)|^2 + (\lambda + \mu) |\operatorname{div}(v - u)|^2) dx + \int_{\Omega} |\partial_t(c - z)|^2 dx \\
 & = - \int_{\Omega} p(\rho|r) \operatorname{div} u dx - \int_{\Omega} \psi''(r)(\rho - r)g dx - \int_{\Omega} h \partial_t(c - z) dx \\
 & - \int_{\Omega} \nabla(c - z) \cdot ((\rho - r)u) dx + \int_{\Omega} (c - z)g dx \\
 & - \int_{\Omega} \rho(v - u) \otimes (v - u) : \nabla u dx - \frac{1}{\zeta} \int_{\Omega} \rho |v - u|^2 dx \\
 & - \int_{\Omega} \left( \frac{\rho - r}{r} (\mu \Delta u + (\lambda + \mu) \nabla \operatorname{div} u) + \rho f \right) \cdot (v - u) dx.
 \end{aligned}$$

This shows (10) and finishes the proof. □

### 4. Weak–strong Uniqueness

We split the proof in several steps.

*Step 1: Relative energy inequality.* We claim that (10) holds for finite energy weak solutions  $(\rho, v, c)$  and  $(\bar{\rho}, \bar{v}, \bar{c})$ , where  $(\bar{\rho}, \bar{v})$  satisfies the regularity (9). According to [9, Section 4], using a density argument, the relative energy inequality (10) still holds for functions  $(r, u)$  satisfying the following regularity conditions:

$$\begin{aligned}
 & r \in C^0_{\text{weak}}([0, T]; L^\gamma(\Omega)), \quad u \in C^0_{\text{weak}}([0, T]; L^{2\gamma/(\gamma-1)}(\Omega; \mathbb{R}^3)), \\
 & |\nabla u| \in L^1(0, T; L^\infty(\Omega)) \cap L^2(\Omega \times (0, T)), \quad u = 0 \text{ on } \partial\Omega, \\
 & \partial_t u \in L^1(0, T; L^{2\gamma/(\gamma-1)}(\Omega; \mathbb{R}^3)) \cap L^2(0, T; L^{6\gamma/(5\gamma-6)}(\Omega; \mathbb{R}^3)), \\
 & |\nabla^2 u| \in L^1(0, T; L^{2\gamma/(2\gamma+1)}(\Omega)) \cap L^2(0, T; L^{6/5}(\Omega)).
 \end{aligned} \tag{36}$$

Moreover,  $r$  needs to be bounded away from zero and we require  $\nabla \psi'(r), \partial_t \psi'(r) \in L^1(0, T; L^{2\gamma/(\gamma-1)}(\Omega))$ . An inspection of (10) reveals that  $z$  should satisfy

$$\begin{aligned}
 & z \in C^0_{\text{weak}}([0, T]; H^1(\Omega)) \cap H^1(0, T; L^2(\Omega)), \quad \Delta z \in L^2(0, T; L^2(\Omega)), \\
 & |\nabla z| \in L^1(0, T; L^{2\gamma/(2\gamma-1)}(\Omega)) \cap L^2(0, T; L^{6\gamma/(5\gamma-6)}(\Omega)).
 \end{aligned} \tag{37}$$

It follows from [7, Theorem 2.4] that (10) still holds if  $(\rho, v, c)$  is a finite energy weak solution.

**Lemma 8.** *Let  $(\bar{\rho}, \bar{v}, \bar{c})$  be a finite energy weak solution in the sense of Definition 1 satisfying the additional regularity (9). Furthermore, let  $\bar{c}^0 \in W^{2-2/\gamma, \gamma}(\Omega)$  and  $\bar{c}^0 \geq 0$  in  $\Omega$ . Then  $(\bar{\rho}, \bar{v}, \bar{c})$  fulfills the regularity conditions (36)–(37).*

*Proof.* Regularity (36) follows as in [9, Section 4] from Sobolev embeddings. Theorem 10 in the Appendix shows that (37) is satisfied.  $\square$

The previous lemma shows that we can take  $(r, u, z) = (\bar{\rho}, \bar{v}, \bar{c})$  in (10). Then the remainder  $R(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c})$  in Lemma 7 simplifies, since  $f = 0$  and  $g = h = 0$ , and we find that

$$\begin{aligned} \int_0^t R(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c})ds &= J_1 + \dots + J_5, \quad \text{where} \\ J_1 &= - \int_0^t \int_{\Omega} p(\rho|\bar{\rho}) \operatorname{div} \bar{v} dx ds, \\ J_2 &= - \int_0^t \int_{\Omega} \nabla(c - \bar{c}) \cdot ((\rho - \bar{\rho})\bar{v}) dx ds, \\ J_3 &= - \int_0^t \int_{\Omega} \rho(v - \bar{v}) \otimes (v - \bar{v}) : \nabla \bar{v} dx ds, \\ J_4 &= - \frac{1}{\zeta} \int_0^t \int_{\Omega} \bar{\rho} |v - \bar{v}|^2 dx ds, \\ J_5 &= - \int_0^t \int_{\Omega} \frac{\rho - \bar{\rho}}{\bar{\rho}} (\mu \Delta \bar{v} + (\lambda + \mu) \nabla \operatorname{div} \bar{v}) \cdot (v - \bar{v}) dx ds. \end{aligned} \tag{38}$$

*Step 2: Estimation of  $J_i$ .* The terms  $J_i$  can be estimated as in [9, Section 4.1] except the new term  $J_2$ . Indeed, since  $p(\rho) = (\gamma - 1)\psi(\rho)$ , we have  $p(\rho|\bar{\rho}) = (\gamma - 1)\psi(\rho|\bar{\rho})$ , showing that

$$J_1 \leq C \int_0^t \int_{\Omega} \psi(\rho|\bar{\rho}) dx ds,$$

and Hölder’s inequality gives

$$J_3 \leq C \int_0^t \int_{\Omega} \rho |v - \bar{v}|^2 dx ds,$$

where  $C > 0$  depends on the  $L^\infty(\Omega \times (0, T))$  norm of  $\nabla \bar{v}$ . The term  $J_4$  is nonpositive and can be neglected. Formulas (4.13)–(4.14) in [9] lead to

$$J_5 \leq \xi \int_0^t \|v - \bar{v}\|_{H^1(\Omega)}^2 ds + C(\xi) \int_0^t \int_{\Omega} \psi(\rho|\bar{\rho}) dx ds,$$

where  $\xi > 0$  is arbitrary and  $C(\xi) > 0$  depends on  $\xi$  as well as  $\|\bar{v}\|_{L^\infty(0,t;W^{2,3}(\Omega))}$  and  $\|\nabla^2 \bar{v}\|_{L^\infty(0,t;L^q(\Omega))}$ . At this point, we need the condition  $q > 3$ .

To estimate the term  $J_2$ , which is not contained in [9], we use equation (3) for  $c$  and integrate by parts:

$$\begin{aligned} J_2 &= - \int_0^t \int_{\Omega} \nabla(c - \bar{c}) \cdot \bar{v} (\partial_t(c - \bar{c}) - \Delta(c - \bar{c}) + (c - \bar{c})) dx \\ &= - \int_0^t \int_{\Omega} \partial_t(c - \bar{c}) \nabla(c - \bar{c}) \cdot \bar{v} dx ds - \frac{1}{2} \int_0^t \int_{\Omega} \nabla[(c - \bar{c})^2] \cdot \bar{v} dx ds \\ &\quad + \int_0^t \int_{\Omega} \left( \operatorname{div} (\nabla(c - \bar{c}) \otimes \nabla(c - \bar{c})) - \frac{1}{2} \nabla |\nabla(c - \bar{c})|^2 \right) \cdot \bar{v} dx ds \\ &= - \int_0^t \int_{\Omega} \partial_t(c - \bar{c}) \nabla(c - \bar{c}) \cdot \bar{v} dx ds + \frac{1}{2} \int_0^t \int_{\Omega} (c - \bar{c})^2 \operatorname{div} \bar{v} dx ds \end{aligned}$$

$$- \int_0^t \int_{\Omega} \left( \nabla(c - \bar{c}) \otimes (c - \bar{c}) : \nabla \bar{v} - \frac{1}{2} |\nabla(c - \bar{c})|^2 \operatorname{div} \bar{v} \right) dx ds.$$

Then, by Young’s inequality,

$$J_2 \leq \frac{1}{2} \int_0^t \int_{\Omega} |\partial_s(c - \bar{c})|^2 dx ds + C \int_0^t \int_{\Omega} (|\nabla(c - \bar{c})|^2 + (c - \bar{c})^2) dx ds,$$

where  $C > 0$  depends on  $\|\bar{v}\|_{L^\infty(0,T;W^{1,\infty}(\Omega))}$ . Summarizing, it follows from (38) that

$$\begin{aligned} \int_0^t R(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) ds &\leq \frac{1}{2} \int_0^t \int_{\Omega} |\partial_s(c - \bar{c})|^2 dx ds + \xi \int_0^t \|v - \bar{v}\|_{H^1(\Omega)}^2 ds \\ &+ C \int_0^t \int_{\Omega} (\psi(\rho|\bar{\rho}) + \rho|v - \bar{v}|^2 + |\nabla(c - \bar{c})|^2 + (c - \bar{c})^2) dx ds. \end{aligned}$$

The first term on the right-hand side can be absorbed by the last term on the left-hand side of (10). The second term on the left-hand side of (10) can be bounded from below by Korn’s inequality [15, Lemma 2] according to

$$\int_{\Omega} (\mu|\nabla(v - \bar{v})|^2 + (\lambda + \mu)|\operatorname{div}(v - \bar{v})|^2) dx \geq C_K \|v - \bar{v}\|_{H^1(\Omega)}^2,$$

since  $v = \bar{v} = 0$  on  $\partial\Omega$ . Therefore, choosing  $0 < \xi < C_K$ , (10) yields

$$\begin{aligned} E((\rho, v, c)(t)|(\bar{\rho}, \bar{v}, \bar{c})(t)) &+ \frac{1}{2} \int_0^t \int_{\Omega} |\partial_s(c - \bar{c})|^2 dx ds + (C_K - \xi) \int_0^t \|v - \bar{v}\|_{H^1(\Omega)}^2 ds \\ &\leq E(\rho_0, v_0, c_0|\bar{\rho}_0, \bar{v}_0, \bar{c}_0) \\ &+ C \int_0^t \int_{\Omega} (\psi(\rho|\bar{\rho}) + \rho|v - \bar{v}|^2 + |\nabla(c - \bar{c})|^2 + (c - \bar{c})^2) dx ds \\ &\leq E(\rho_0, v_0, c_0|\bar{\rho}_0, \bar{v}_0, \bar{c}_0) + C \int_0^t H(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) ds. \end{aligned} \tag{39}$$

*Step 3: Estimation of  $\int_{\Omega} (\rho - \bar{\rho})(c - \bar{c}) dx$ .* We use Lemma 9 in Appendix 4 with  $m = 2$  and arbitrary  $\kappa_1, \xi > 0$  on the set  $\{\rho \leq R\}$  for some  $R > 0$ :

$$\begin{aligned} \int_{\{\rho \leq R\}} (\rho - \bar{\rho})(c - \bar{c}) dx &\leq \kappa_1 \|\rho - \bar{\rho}\|_{L^2(\Omega \cap \{\rho \leq R\})}^2 + \xi \|\nabla(c - \bar{c})\|_{L^2(\Omega)}^2 \\ &+ C_1(\kappa_1, \xi) \|c - \bar{c}\|_{L^1(\Omega)}^{C_2(2)}, \end{aligned} \tag{40}$$

as well as with  $m = \gamma$  (which requires  $\gamma > 8/5$ ) and arbitrary  $\kappa_2 > 0$  on the set  $\{\rho > R\}$ :

$$\begin{aligned} \int_{\{\rho > R\}} (\rho - \bar{\rho})(c - \bar{c}) dx &\leq \kappa_2 \|\rho - \bar{\rho}\|_{L^\gamma(\Omega \cap \{\rho > R\})}^\gamma + \xi \|\nabla(c - \bar{c})\|_{L^2(\Omega)}^2 \\ &+ C_1(\kappa_2, \xi) \|c - \bar{c}\|_{L^1(\Omega)}^{C_2(\gamma)}. \end{aligned} \tag{41}$$

According to [13, Lemma 2.4], there exist constants  $C_3, C_4, c_p, C_p > 0$  such that

$$\psi(\rho|\bar{\rho}) \geq \begin{cases} C_3|\rho - \bar{\rho}|^2 & \text{if } 0 \leq \rho \leq R, \\ C_4|\rho - \bar{\rho}|^\gamma & \text{if } \rho > R, \end{cases}$$

as long as  $c_p \leq \bar{\rho} \leq C_p$ . Thus, we can replace the first term on the right-hand sides of (40) and (41), respectively, by  $\kappa_1 C_3^{-1} \int_{\Omega} \psi(\rho|\bar{\rho}) dx$  and  $\kappa_2 C_4^{-1} \int_{\Omega} \psi(\rho|\bar{\rho}) dx$ , and summing these inequalities, we obtain

$$\begin{aligned} \int_{\Omega} (\rho - \bar{\rho})(c - \bar{c}) dx &\leq \left( \frac{\kappa_1}{C_3} + \frac{\kappa_2}{C_4} \right) \int_{\Omega} \psi(\rho|\bar{\rho}) dx + 2\xi \|\nabla(c - \bar{c})\|_{L^2(\Omega)}^2 \\ &+ C_1(\kappa_1, \xi) \|c - \bar{c}\|_{L^1(\Omega)}^{C_2(2)} + C_1(\kappa_2, \xi) \|c - \bar{c}\|_{L^1(\Omega)}^{C_2(\gamma)}. \end{aligned} \tag{42}$$



We wish to estimate the last two norms in terms of the initial data. To this end, we integrate (3) and use the mass conservation  $\|\rho(t)\|_{L^1(\Omega)} = \|\rho^0\|_{L^1(\Omega)}$  and  $\|\bar{\rho}(t)\|_{L^1(\Omega)} = \|\bar{\rho}^0\|_{L^1(\Omega)}$ :

$$\frac{d}{dt} \int_{\Omega} (c - \bar{c})(t) dx = - \int_{\Omega} (c - \bar{c}) dx + \int_{\Omega} (\rho - \bar{\rho}) dx = - \int_{\Omega} (c - \bar{c}) dx + \int_{\Omega} (\rho^0 - \bar{\rho}^0) dx.$$

Gronwall’s lemma yields

$$\int_{\Omega} (c - \bar{c})(t) dx \leq C \int_{\Omega} (c^0 - \bar{c}^0) dx + C \int_{\Omega} (\rho^0 - \bar{\rho}^0) dx.$$

The same argument with  $\bar{c} - c$  then shows that

$$\|(c - \bar{c})(t)\|_{L^1(\Omega)} \leq C (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)}).$$

Hence, choosing  $\kappa_1 = C_3/4$ ,  $\kappa_2 = C_4/4$ , and  $\xi = 1/8$ , we deduce from (42) that

$$\begin{aligned} \int_{\Omega} (\rho - \bar{\rho})(c - \bar{c}) dx &\leq \frac{1}{2} \int_{\Omega} \left( \psi(\rho|\bar{\rho}) + \frac{1}{2} |\nabla(c - \bar{c})|^2 \right) dx \\ &\quad + C_1(\kappa_1, \xi) (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)})^{C_2(2)} \\ &\quad + C_1(\kappa_2, \xi) (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)})^{C_2(\gamma)}. \end{aligned}$$

The last two terms are bounded from above by

$$C (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)})^{C_5},$$

where  $C_5$  equals  $C_2(2)$  or  $C_2(\gamma)$  depending on whether  $\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)}$  is smaller or larger than one. We conclude that

$$\begin{aligned} \int_{\Omega} (\rho - \bar{\rho})(c - \bar{c}) dx &\leq \frac{1}{2} \int_{\Omega} \left( \psi(\rho|\bar{\rho}) + \frac{1}{2} |\nabla(c - \bar{c})|^2 \right) dx \\ &\quad + C (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)})^{C_5} \\ &\leq \frac{1}{2} H(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) + C (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)})^{C_5}. \end{aligned} \tag{43}$$

*Step 4: End of the proof.* By (43), the relative energy is bounded from below by

$$\begin{aligned} E(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) &\geq H(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) - \int_{\Omega} (\rho - \bar{\rho})(c - \bar{c}) dx \\ &\geq \frac{1}{2} H(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) - C (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)})^{C_5}. \end{aligned}$$

We insert this estimate into (39):

$$\begin{aligned} &\frac{1}{2} H((\rho, v, c)(t)|(\bar{\rho}, \bar{v}, \bar{c})(t)) + \frac{1}{2} \int_0^t \int_{\Omega} |\partial_s(c - \bar{c})|^2 dx ds + C \int_0^t \|v - \bar{v}\|_{H^1(\Omega)}^2 ds \\ &\leq E(\rho^0, v^0, c^0|\bar{\rho}^0, \bar{v}^0, \bar{c}^0) + C \int_0^t H(\rho, v, c|\bar{\rho}, \bar{v}, \bar{c}) ds \\ &\quad + C (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)})^{C_5}. \end{aligned}$$

An application of Gronwall’s lemma gives

$$\begin{aligned} H((\rho, v, c)(t)|(\bar{\rho}, \bar{v}, \bar{c})(t)) &\leq C e^{Ct} \{ E(\rho^0, v^0, c^0|\bar{\rho}^0, \bar{v}^0, \bar{c}^0) \\ &\quad + (\|c^0 - \bar{c}^0\|_{L^1(\Omega)} + \|\rho^0 - \bar{\rho}^0\|_{L^1(\Omega)})^{C_5} \}, \end{aligned}$$

and the choice  $\rho^0 = \bar{\rho}^0$ ,  $v^0 = \bar{v}^0$ ,  $c^0 = \bar{c}^0$  ends the proof.

**Funding** Open access funding provided by Austrian Science Fund (FWF).

**Conflict of interest** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Open Access.** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

### Appendix A: Auxiliary Results

**Lemma 9.** *Let  $\Omega \subset \mathbb{R}^d$  be a bounded domain with  $d \in \{2, 3\}$  and let  $m > 2(d+1)/(d+2)$ . Furthermore, let  $\kappa, \xi > 0$ . Then there exist constants  $C_1(\kappa, \xi) > 0$  and  $C_2(m) > 0$  such that for all  $\rho \in L^m(\Omega)$ ,  $c \in H^1(\Omega)$ ,*

$$\int_{\Omega} \rho c dx \leq \kappa \|\rho\|_{L^m(\Omega)}^m + \xi \|\nabla c\|_{L^2(\Omega)}^2 + C_1(\kappa, \xi) \|c\|_{L^1(\Omega)}^{C_2(m)}.$$

*Proof.* The proof of the lemma is contained in [18, Appendix B] for solutions to the degenerate Keller–Segel equations. For clarity, we present the proof for general functions  $\rho$  and  $c$ . We conclude from the interpolation inequality for Lebesgue spaces and Young’s inequality that for any  $\kappa > 0$ ,

$$\|\rho c\|_{L^1(\Omega)} \leq \|\rho\|_{L^m(\Omega)} \|c\|_{L^{m/(m-1)}(\Omega)} \leq \kappa \|\rho\|_{L^m(\Omega)}^m + C(\kappa) \|c\|_{L^{m/(m-1)}(\Omega)}^{m/(m-1)}.$$

We estimate the second term on the right-hand side by applying the Gagliardo–Nirenberg inequality with  $\theta = 2d/(m(d+2))$ :

$$\|c\|_{L^{m/(m-1)}(\Omega)} \leq C \|\nabla c\|_{L^2(\Omega)}^{\theta} \|c\|_{L^1(\Omega)}^{1-\theta} + C \|c\|_{L^1(\Omega)}.$$

Then, by Minkowski’s and Young’s inequality, for any  $\varepsilon > 0$ ,

$$\begin{aligned} \|\rho c\|_{L^1(\Omega)} &\leq \kappa \|\rho\|_{L^m(\Omega)}^m + C(\kappa, m) (\|\nabla c\|_{L^2(\Omega)}^{m\theta/(m-1)} \|c\|_{L^1(\Omega)}^{m(1-\theta)/(m-1)} + \|c\|_{L^1(\Omega)}^{m/(m-1)}) \\ &\leq \kappa \|\rho\|_{L^m(\Omega)}^m + C(\kappa, m) \varepsilon \|\nabla c\|_{L^2(\Omega)}^2 \\ &\quad + C(\kappa, m, \varepsilon) (\|c\|_{L^1(\Omega)}^{2m(1-\theta)/(2(m-1)-m\theta)} + \|c\|_{L^1(\Omega)}^{m/(m-1)}), \end{aligned}$$

which is possible since  $m\theta/(m-1) < 2$  is equivalent to  $m > 2(d+1)/(d+2)$ . The lemma follows after choosing  $\varepsilon = \xi/C(\kappa, m)$ ,  $C_1(\kappa, \xi) = C(\kappa, m, \varepsilon)$ , and  $C_2(m) = \max\{m/(m-1), 2m(1-\theta)/(2(m-1)-m\theta)\}$ .  $\square$

The following result concerns the maximal regularity of the solution to

$$\partial_t u - \Delta u + u = f \quad \text{in } \Omega, \quad t > 0, \tag{44}$$

$$\nabla u \cdot \nu = 0 \text{ on } \partial\Omega, \quad t > 0, \quad u(\cdot, 0) = u^0 \text{ in } \Omega, \tag{45}$$

where  $\Omega \subset \mathbb{R}^d$  ( $d \geq 1$ ) is a bounded domain with  $C^3$  boundary. We recall that  $W_{\nu}^{2,-2/p,q}(\Omega)$  is the completion of the space of functions  $w \in C^{\infty}(\bar{\Omega})$  satisfying  $\nabla w \cdot \nu = 0$  on  $\partial\Omega$  in the norm of  $W^{2,-2/p,q}(\Omega)$ . The theorem is a special case of [6, Theorem 10.22] or [16, Lemma 7.37].

**Theorem 10** (*Maximal regularity*). *Let  $1 < p, q < \infty$ ,  $f \in L^p(0, T; L^q(\Omega))$ , and let  $u^0 \in W_{\nu}^{2,-2/p,q}(\Omega)$ . Then there exists a unique solution  $u$  to (44)–(45) satisfying*

$$u \in L^p(0, T; W^{2,q}(\Omega)) \cap W^{1,p}(0, T; L^q(\Omega)) \cap C^0([0, T]; W^{2,-2/p,q}(\Omega)),$$

and there exists a constant  $C > 0$  such that

$$\begin{aligned} & \|u\|_{L^\infty(0,T;W^{2-2/p,q}(\Omega))} + \|u\|_{L^p(0,T;W^{2,q}(\Omega))} + \|\partial_t u\|_{L^p(0,T;L^q(\Omega))} \\ & \leq C(\|f\|_{L^p(0,T;L^q(\Omega))} + \|u^0\|_{W^{2-2/p,q}(\Omega)}). \end{aligned}$$

## References

- [1] Aïssa, N., Alexandre, R.: Global existence of weak solutions to an angiogenesis model. *J. Evol. Eqs.* **16**, 877–894 (2016)
- [2] Ambrosi, D., Gamba, A., Serini, G.: Cell directional and chemotaxis in vascular morphogenesis. *Bull. Math. Biol.* **66**, 1851–1873 (2004)
- [3] Biler, P., Brandolese, L.: On the parabolic-elliptic limit of the doubly parabolic Keller–Segel system modelling chemotaxis. *Studia Math.* **193**, 241–261 (2009)
- [4] Di Francesco, M., Donatelli, D.: Singular convergence of nonlinear hyperbolic chemotaxis systems to Keller–Segel type models. *Discrete Cont. Dyn. Sys. B* **13**, 79–100 (2010)
- [5] Feireisl, E.: *Dynamics of Viscous Compressible Fluids*. Oxford University Press, Oxford (2004)
- [6] Feireisl, E., Novotný, A.: *Singular Limits in Thermodynamics of Viscous Fluids*. Birkhäuser, Basel (2009)
- [7] Feireisl, E., Jin, B.J., Novotný, A.: Relative entropies, suitable weak solutions, and weak–strong uniqueness for the compressible Navier–Stokes system. *J. Math. Fluid Mech.* **14**, 717–730 (2012)
- [8] Feireisl, E., Novotný, A., Petzeltová, H.: On the existence of globally defined weak solutions to the Navier–Stokes equations. *J. Math. Fluid Mech.* **3**, 358–392 (2001)
- [9] Feireisl, E., Novotný, A., Sun, Y.: Suitable weak solutions to the Navier–Stokes equations of compressible viscous fluids. *Indiana Univ. Math. J.* **60**, 611–631 (2011)
- [10] Germain, P.: Weak-strong uniqueness for the isentropic compressible Navier–Stokes system. *J. Math. Fluid Mech.* **13**, 137–146 (2011)
- [11] Huo, X., Jüngel, A., Tzavaras, A.E.: Weak-strong uniqueness for Maxwell–Stefan systems. *SIAM J. Math. Anal.* **54**, 3215–3252 (2022)
- [12] Jiang, S., Zhang, P.: Axisymmetric solutions of the 3D Navier–Stokes equations for compressible isentropic fluids. *J. Math. Pures Appl.* **82**, 949–973 (2003)
- [13] Lattanzio, C., Tzavaras, A.: Relative entropy in diffusive relaxation. *SIAM J. Math. Anal.* **45**, 1563–1584 (2013)
- [14] Mellet, A., Vasseur, A.: On the barotropic compressible Navier–Stokes equations. *Commun. Partial Differ. Eqs.* **32**, 431–452 (2007)
- [15] Novotný, A., Pokorný, M.: Steady compressible Navier–Stokes–Fourier system for monoatomic gas and its generalizations. *J. Differ. Eqs.* **251**, 270–315 (2011)
- [16] Novotný, A., Straškraba, I.: *Introduction to the Mathematical Theory of Compressible Flow*. Oxford University Press, Oxford (2004)
- [17] Serini, G., Ambrosi, D., Girauso, E., Gamba, A., Preziosi, L., Bussolini, F.: Modeling the early stages of vascular network assembly. *EMBO J.* **22**, 1771–1779 (2003)
- [18] Sugiyama, Y.: Time global existence and asymptotic behavior of solutions to degenerate quasi-linear parabolic systems of chemotaxis. *Differ. Integral Eqs.* **20**, 133–180 (2007)
- [19] Tao, Y., Winkler, M.: Boundedness in a quasilinear parabolic-parabolic Keller–Segel system with subcritical sensitivity. *J. Differ. Eqs.* **252**, 692–715 (2012)
- [20] Tosin, A., Ambrosi, D., Preziosi, L.: Mechanics and chemotaxis in the morphogenesis of vascular networks. *Bull. Math. Biol.* **68**, 1819–1836 (2006)
- [21] Tuval, I., Cisneros, L., Dombrowski, C., Wolgemuth, C., Kessler, J., Goldstein, R.: Bacterial swimming and oxygen transport near contact lines. *PNAS* **102**, 2277–2282 (2005)
- [22] Winkler, M.: Global large-data solutions in a chemotaxis– (Navier)–Stokes system modeling cellular swimming in fluid drops. *Commun. Partial Differ. Eqs.* **37**, 319–351 (2012)

Xiaokai Huo  
 Department of Mathematics  
 Iowa State University  
 411 Morrill Road  
 Ames IA 50011-2104  
 USA  
 e-mail: xhuo@iastate.edu

Ansgar Jüngel  
 Institute of Analysis and Scientific Computing  
 Technische Universität Wien  
 Wiedner Hauptstrasse 8–10  
 1040 Wien  
 Austria  
 e-mail: juengel@tuwien.ac.at

(accepted: October 19, 2023)