

ROLE OF SIALIC ACID AND GLYCOSYLATIONS AS BIOMARKERS FOR SEVERITY STRATIFICATION IN POST-VIRAL INFECTION SYNDROME


Davide Ret^{a,b}, Daniel García de Otazo Hernández^{a,b}, Gianluigi Sabatino^a, Selina Keppler^c, Eva Untersmayr^b

^aResearch Unit Macromolecular Chemistry, Institute of Applied Synthetic Chemistry, TU Wien, 1060 Vienna, Austria, davide.ret@tuwien.ac.at

^bInstitute of Pathophysiology and Allergy Research, Center for Pathophysiology, Infectiology and Immunology, Medical University of Vienna, 1090 Vienna, Austria

^cDivision of rheumatology and Immunology Medical University of Graz, 8010 Graz, Austria

Our group focuses on the study of sialic acid and sialylated glycans in inflammatory diseases. In this work, we aim to find better biomarkers for severity stratification in post-viral syndrome using a glycomics approach [1]. Glycomics is the communication language of organisms, based on a sugar code (glycans) imprinted on tissues, proteins, and lipids. Many diseases, such as autoimmune disorders and pathogenic infections, has an altered sugar code in the organism, leading to inflammation.



N-glycome was characterized in serum and purified IgG by MALDI mass spectrometry in positive mode after purification and derivatization with methylamine [2]. After labelling with DMBA, total sialic acid content was quantified by HPLC-RP-FL.

We observed variations in N-glycome and sialic acid concentration between patients and controls, suggesting the involvement of sialic acid in the severity of post-viral fatigue syndrome.

1. Rohrhofer, J., et al., *Immunological Patient Stratification in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome*. Journal of Clinical Medicine, **2024**. 13(1): p. 275.
2. Ret, D., et al., *DMTMM-mediated methylamidation for MALDI mass spectrometry analysis of N-glycans with structurally conserved sialic acid residues in biological fluids "via direttissima"*. Talanta, **2022**. 242: p. 123326.